

OPEN

A new possible megalosauroid theropod from the Middle Jurassic Xintiangou Formation of Chongqing, People's Republic of China and its implication for early tetanuran evolution

Hui Dai^{1*}, Roger Benson², Xufeng Hu¹, Qingyu Ma¹, Chao Tan¹, Ning Li¹, Ming Xiao¹, Haiqian Hu¹, Yuxuan Zhou¹, Zhaoying Wei¹, Feng Zhang¹, Shan Jiang³, Deliang Li⁴, Guangzhao Peng³, Yilun Yu^{5,6} & Xing Xu^{5,6*}

Tetanurae is a special group of theropod dinosaurs that originated by the late Early Jurassic. It includes several early-diverging groups of generally large-bodied predators (megalosauroids, allosauroids, tyrannosauroid coelurosaurs) as well as morphologically disparate small-bodied coelurosaurs, including birds. Aspects of the evolutionary history of tetanurans remain contested, including the topology of their deep phylogenetic divergences (among Megalosauroida, Allosauroida and Coelurosauria). We report a new theropod, *Yunyangosaurus puanensis* gen. et sp. nov., based on a fragmentary specimen recovered from the Middle Jurassic Xintiangou Formation of Chongqing, southwestern China. It shares several features uniquely with some megalosauroids (the clade of megalosauroids + spinosaurids + piatnitzkysaurids), such as prominent rims around the anterior articular surfaces of cervical centra and bifurcated anterior dorsal neural spines (present in piatnitzkysaurids). Nevertheless, it also shows several features that are rare or absent among megalosauroids and more crownward tetanurans, including prominent spinopostzygopophyseal laminae (also present in non-tetanurans and metriacanthosaurid allosauroids), flat anterior articular surfaces of the cervical centra (also present in piatnitzkysaurids and some earlier-diverging tetanurans), and the presence of a posterior pneumatic foramen or fossa (absent in most tetanurans, but sporadically present in some cervical vertebrae of piatnitzkysaurids). *Yunyangosaurus* therefore presents a combination of derived and apparently primitive character states that are not seen in other theropods. This suggests that patterns of morphological evolution associated with deep tetanuran divergences were more complex than currently recognized, with implications for understanding the character evolution in theropods.

The Middle and Upper Jurassic Shaximiao Formation of southwestern China has produced spectacular fossil remains of dinosaurs and many other vertebrates^{1,2}. Significant discoveries from this formation in the Chongqing area include the sauropods *Mamenchisaurus hochuanensis*, *Omeisaurus changshouensis*, and the theropod *Yangchuanosaurus shangyouensis*³ as well as the stegosaur *Chungkingosaurus jiangbeiensis*⁴. Compared to the

¹Chongqing Laboratory of Geoheritage Protection and Research, No. 208 Hydrogeological and Engineering Geological Team, Chongqing Bureau of Geological and Mineral Resource Exploration and Development, Chongqing, 400700, China. ²Department of Earth Sciences, University of Oxford, Oxford, OX1 3AN, UK. ³Zigong Dinosaur Museum, Zigong, 643013, Sichuan, China. ⁴Chongqing Institute of Geological Survey, Chongqing, 401122, China. ⁵Key Laboratory of Vertebrate Evolution and Human Origins of Chinese Academy of Sciences, Institute of Vertebrate Paleontology and Paleoanthropology, Chinese Academy of Sciences, Beijing, 100044, China. ⁶CAS Center for Excellence in Life and Palaeoenvironment, Beijing, China. *email: dhui6251@qq.com; xu.xing@ivpp.ac.cn




Figure 1. Geographical map indicating the location of the fossil site within the Yunyang County, Chongqing Municipality, China. Modified from Li.

fossiliferous Shaximiao Formation, the underlying lower Middle Jurassic Xintiangou Formation^{5,6} has produced few vertebrate fossils. In 2016, we organized a survey in the upper portion of the Xintiangou Formation of Laojun Village, Puan Township, Yunyang City, Chongqing (Fig. 1). This resulted in the discoveries of numerous vertebrate fossils, including a fragmentary theropod specimen recovered from a layer of gray lamellar shell siltstone intercalated with occasional silty mudstone (Fig. 2). The specimen comprises only some presacral vertebrae and several fragmentary bones, but it displays some informative features for its systematic position and provides new information on the early evolution of tetanuran dinosaurs. In the present paper, we establish a new tetanuran species based on this specimen, describe the specimen, and present a short discussion on its implications for the evolution of anatomical character states among early tetanurans.

Results

Systematic palaeontology

Theropoda Marsh 1881

Tetanurae Gauthier 1986

Yunyangosaurus puanensis gen. et sp. nov.

Etymology. Generic name is a combination of Yunyang (the county-level city in which the type locality is located) and sauros (Greek, reptile). The specific name is derived from Puan (the town name of the type locality).

Holotype. CLGPR v00002 (Chongqing Laboratory of Geoheritage Protection and Research), a partial skeleton comprising eleven presacral vertebrae, several cervical and dorsal ribs and chevrons. These bones are disarticulated, but were found associated in a small area about 5 square metres, with no other theropod skeletal elements preserved nearby.

Locality and horizon. Laojun Village, Puan Township, Yunyang City, Chongqing, China (Fig. 2). Middle Jurassic Xintiangou Formation^{5,6}. It should be noted that a recent U-Pb dating of the lowest beds of the overlying Shaximiao Formation suggested that the Shaximiao Formation started deposition in late Oxfordian⁷, but that study has not been widely accepted⁵. Furthermore, our newly collected unpublished U-Pb dating data support the traditional view that lower beds of Shaximiao Formation is Middle Jurassic in age.

Diagnosis. An early-branching tetanuran distinguishable from other tetanurans by the following unique combination of features (*indicates autapomorphies): axial intercentrum sub-triangular in lateral view (sub-rectangular in *Allosaurus* and *Sinraptor*); mediolateral width of axial centrum tapers ventrally, resulting in a subtriangular outline in posterior view*; axis with an accessory medial spinopostzygapophyseal lamina* and a ball-like structure at the base of anterior margin of the axial neural spine*; axis bears very prominent spinopostzygapophyseal lamina similar to non-tetanuran theropods and metriacanthosaurids such as *Sinraptor*, but different from most tetanurans, including other Chinese Jurassic tetanurans and patnitzyksaurids; cervical




Figure 2. Stratigraphic section of the sedimentary sequence at the type locality. Modified from Li.

centra with flat anterior articular surfaces, similar to non-tetanurans, *Condorraptor*, *Piatnitzkysaurus* and some Jurassic Chinese tetanurans such as ‘*Szechuanosaurus*’ *zigongensis* and *Xuanhanosaurus*, but different from most other non-coelurosaurian tetanurans, which have convex anterior surfaces, including *Leshansaurus*, *Marshosaurus* *Monolophosaurus* and metriacanthosaurids such as *Sinraptor*; ventral ridge present in anterior cervicals (absent in most tetanurans, but present in some non-tetanurans and the piatnitzkysaurid *Marshosaurus* as a low, anterior eminence); posterior pneumatic foramen or fossa present in some cervical centra (also present in some of the pneumatic centra of *Piatnitzkysaurus* and *Condorraptor* and in all pneumatic centra of most non-tetanuran theropods; but absent in most tetanurans, including Chinese Jurassic tetanurans *Gasosaurus*, *Leshansaurus*, *Sinraptor*, ‘*Szechuanosaurus*’ *zigongensis*, *Xuanhanosaurus* and *Yangchuanosaurus*); prominent epiphyses present in posterior cervicals (absent or relatively low in other Chinese Jurassic tetanurans where known, such as *Monolophosaurus*, *Sinraptor* and ‘*Szechuanosaurus*’ *zigongensis*, as well as *Carnotaurus*, *Torvosaurus*, *Condorraptor* and *Piatnitzkysaurus*); cervical epiphyses prominent and posteriorly oriented* (posterolaterally oriented in most other theropods); posterior cervical and anterior dorsal vertebrae with transversely bifurcated neural spines*, especially prominent in anterior dorsals (less prominent bifurcated neural

Vert. No.	CL	PH	PW	MW	TH	NSH	NSL	NSW
PV1	58	39	47	14	113	56	34	16
PV 2	69	48	50	15	102+	42+	32	8
PV 3	68	57	64	27	—	—	—	—
PV4	54	54	55+	24	—	—	—	—
PV5	70	59	68	31	106	53	17	10
PV6	58	62	70	29	178	98	20	18
PV7	61	61	68	30	168	85	20	14
PV8	61	54	52+	36	—	—	—	—
PV9	52	61	54+	25	164	90	18	17
PV10	60	60	65	32	—	78+	30	18
PV11	63	57	67	28	151+	71+	37	14

Table 1. Vertebral measurements (in millimetres) of *Yunyangosaurus puanensis* holotype. The vertebrae are named as PV1 to PV11 for the convenience of the description; Abbreviations: CL, centrum length; PH, centrum posterior end height; PW, centrum posterior end width; MW, centrum middle portion width; TH, vertebra total height (measured at posterior end); NSH, neural spine height; NSL, neural spine anteroposterior length; NSW, neural spine width; +, incomplete measurements.

spines are present in anterior dorsals only in *Condoraptor*, *Marshosaurus* and *Piatnitzkysaurus*; but absent in Chinese Jurassic theropods where preserved, such as *Sinraptor*, *Monolophosaurus*, ‘*Szechuanosaurus*’ *zigongensis* and *Yangchuanosaurus*); prominent hypapophyses in anterior dorsal vertebrae (also seen in metriacanthosaurids, but small in *Monolophosaurus* and most other early-branching tetanurans); distinctive fossae immediately ventral to dorsal parapophyses*.

Description and comparisons. The *Yunyangosaurus puanensis* holotype is inferred to be a sub-adult individual based on the closure of neurocentral sutures in most preserved vertebrae. It has an estimated total body length of 4.7 meters based on the similarity of the measurements of several vertebrae (Table 1) to those of *Sinraptor dongi*⁸. It is therefore similar to species such as *Gasosaurus constructus*⁴ and *Piatnitzkysaurus floresi* from the late Early Jurassic of Argentina⁹. However, it is smaller in body size than most other known early-branching tetanurans from the Jurassic of China^{1,8} with the exception of *Chuangongocoelurus*, the type specimen of which is considerably smaller^{10,11}.

Throughout our description, we aim to make comparisons with Chinese Jurassic tetanurans where possible. Among these taxa, *Sinraptor*, *Yangchuanosaurus* and *Shidaisaurus* are metriacanthosaurids¹⁰ with well-known and relatively similar anatomy to each other^{3,12–15}. In general, we have compared *Yunyangosaurus* to *Sinraptor* as an almost complete representative of this clade. Other Chinese Jurassic tetanurans potentially occupy a wide range of phylogenetic positions among early-diverging tetanurans. However, extensive comparisons with some of these are more difficult. The referred materials of some taxa, such as *Chuangongocoelurus* and *Kaijiangosaurus*, are likely represented by specimens from multiple species¹⁰, so the status of materials that overlap with the preserved material of *Yunyangosaurus* is not clear and requires revision. These taxa are not compared to *Yunyangosaurus* in our description. Nevertheless, they show clear differences: referred vertebral centra of *Chuangongocoelurus* are proportionally elongate, and referred cervical vertebrae of *Kaijiangosaurus* have one pleurocoel on each central side and anteroposteriorly broad and transversely non-bifurcated neural spines¹¹. Other taxa such as *Gasosaurus*, *Leshansaurus*, ‘*Szechuanosaurus*’ *zigongensis* and *Xuanhanosaurus* include overlapping, definitely referable material¹⁶. Of these, vertebrae of *Leshansaurus*, ‘*Szechuanosaurus*’ *zigongensis* are relatively complete and well-preserved¹⁷ whereas *Xuanhanosaurus* includes relatively few, incomplete vertebrae¹⁸ and *Gasosaurus* includes most presacral vertebrae, but they are incomplete and highly reconstructed by plaster, making their original morphology difficult to observe (IVPP)⁴. All four are apparently different from the new taxon, as indicated in our diagnosis and the description below.

Eleven presacral vertebrae of *Yunyangosaurus* are preserved, seven of them are identified as cervical vertebrae, and four as dorsal vertebrae, based on their anatomy (e.g., the position of parapophysis, the orientation of diapophysis, and the presence of a hypapophysis). The cervical series is represented by two anterior cervicals, three middle cervicals, and two posterior cervicals (probably the posteriormost two cervicals). The dorsal series is represented by three anterior dorsal vertebrae (probably the anteriormost ones) and an anterior middle dorsal vertebra.

The axis is mostly preserved, missing the odontoid and part of left spinopostzygapophyseal lamina. The axial intercentrum is attached to the axial centrum, but the line of fusion is obviously visible (Fig. 3A,B). The axial intercentrum is sub-triangular in outline in lateral view. This is similar to the condition in many theropods, including *Leshansaurus*¹⁶, *Marshosaurus* (CMNH 21704), *Monolophosaurus*¹⁹, *Piatnitzkysaurus*⁹, ‘*Szechuanosaurus*’ *zigongensis*¹⁷ and *Neovenator* (Brusatte *et al.*, 2008), but is unlike the sub-rectangular axial intercentra of some allosauroids: *Sinraptor*¹³ and *Allosaurus*²⁰. The ventral surface of the axial intercentrum is horizontal, as in some non-tetanuran theropods, *Piatnitzkysaurus*⁹ and the possible megalosaurid *Leshansaurus*¹⁶. In contrast, the ventral surface of the intercentrum is inclined anterodorsally relative to the ventral surface of the axis, such




Figure 3. Axis (PV1, Table 1) of *Yunyangosaurus puanensis* holotype in anterior (A), left lateral (B), posterior (C), right lateral (D), dorsal (E) and ventral. (F) Axi, axial intercentrum; bs, ball-like structure; dp, diapophysis; ep, epiphysis; ho, hyposphene; nc, neural canal; ns, neural spine; poz, postzygapophysis; pp, parapophysis; prz, prezygapophysis; lspol, lateral spinopostzygapophyseal lamina; mspol, medial spinopostzygapophyseal lamina. Scale bar = 5 cm.

that the two form an oblique angle as in *Monolophosaurus*¹⁹ and some allosauroids such as *Acrocanthosaurus*²¹, *Giganotosaurus* (IVPP V 7265) and *Sinraptor*¹³.

The axial centrum is platycoelous, with a flat anterior surface. The anterior articular surface of the axial centrum faces anterodorsally, as does the articular surface of the axial intercentrum for the atlantal intercentrum. The axial centrum bears a shallow fossa on its lateral surface on each side. These fossae are asymmetrically developed: the right one is large and shallow, but the left one is small and deep (Fig. 3B,D). This fossa, on the left side, is proportionally larger than those of many other tetanurans, including *Allosaurus*²⁰, *Sinraptor*¹³, ‘*Szechuanosaurus*’ *zigongensis*¹⁷, *Eustreptospondylus*²² and *Piatnitzkysaurus* (PVL 4073). The ventral surface of the centrum is transversely narrow with a mediolateral width that tapers ventrally, but nevertheless it is transversely rounded and lacks a distinct ventral keel. This morphology may be an autapomorphy of *Yunyangosaurus* and is unlike the transversely broad ventral surface of the axial centrum in ‘*Szechuanosaurus*’ *zigongensis* (ZDM 9011), *Yangchuanosaurus hepingensis* (ZDM 0024), *Allosaurus*²⁰, *Sinraptor*¹³ and *Eustreptospondylus*²². *Piatnitzkysaurus* (PVL 4073) and *Marshosaurus* (CMNH 21704) have transversely narrow axial centra and are therefore somewhat similar to *Yunyangosaurus*. However, the width of the axial centrum does not taper ventrally in these taxa, unlike in *Yunyangosaurus*. The posterior articular surface of the axial centrum of *Yunyangosaurus* has a sub-triangular outline when seeing in posterior view, and in lateral view it forms an acute angle with the ventral margin (Fig. 3C) to a greater degree than in *Yangchuanosaurus*¹⁸, *Carnotaurus*²³, *Ceratosaurus*²⁴, or *Sinraptor*¹³. The neural canal of the axis is 17 mm high and 18 mm wide while the posterior articular surface is 39 mm high and 47 mm wide, indicating the presence of a relatively large neural canal.

The ventrolaterally oriented axial diapophysis is small and is situated at the fusion line of the centrum and neural arch. The axial prezygapophysis is relatively small and its articular surface faces anterodorsolaterally. The axial postzygapophyses are large with well-developed quadrangular articular surfaces, similar to those of more posterior presacral vertebrae. The articular surfaces face posteroventrally with a lateral inclination of about 20° from horizontal (Fig. 3C). Prominent, broad epiphyses overhang the postzygapophyseal articular surfaces and their long axes are oriented slightly more medially than those of the postzygapophyses. Axial epiphyses are prominent in many non-tetanuran theropods, including *Dilophosaurus*²⁵ and *Carnotaurus*²³, as well as in piatnitzkysaurids (*Piatnitzkysaurus*⁹; *Marshosaurus*, CMNH 21704) and many allosauroids, including metriacanthosauroids such as *Sinraptor*¹³ and *Shidaisaurus*¹⁵, but are somewhat reduced in megalosauroids (*Eustreptospondylus*)^{10,22} and spinosauroids (*Baryonyx*)²⁶.

There are two pairs of spinopostzygapophyseal laminae: a lateral one and a medial one. The prominent lateral one is the major one, and it is comparable to the spinopostzygapophyseal lamina reported in metriacanthosauroids such as *Sinraptor*¹³ and *Shidaisaurus*¹⁵, and in non-tetanuran theropods such as *Ceratosaurus*²⁴ and *Dilophosaurus*²⁵. In contrast, it is either small or absent in most early-branching tetanurans (e.g. *Allosaurus*²⁰; *Baryonyx*²⁶; *Eustreptospondylus*²², *Marshosaurus*, CMNH 21704, *Monolophosaurus*¹⁹ and *Piatnitzkysaurus*⁹). The




Figure 4. An anterior cervical vertebra (PV2, Table 1) of *Yunyangosaurus puanensis* holotype in anterior (A), left lateral (B), posterior (C), right lateral (D), dorsal (E), ventral (F) and 3D images (G,H). cpol centropostzygapophyseal lamina; dp, diapophysis; ep, epiphysis; f, fossa; ipol, intrapostzygapophyseal lamina; nc, neural canal; ns, neural spine; pl, pneumatic foramen (pleurocoel); poz, postzygapophysis; pp, parapophysis; prdl, prezygodipophyseal lamina; prz, prezygapophysis; spl, spinopostzygapophyseal lamina; sprl, spinoprezygapophyseal lamina. Scale bar = 5 cm.

second, more medially located spinopostzygapophyseal lamina, is oriented vertically and is located within the fossa on the posterior surface of the axial neural spine, between the more lateral spinopostzygapophyseal lamina and the central portion of the neural spine. This medial spinopostzygapophyseal lamina has not been reported in other theropods and may be an autapomorphy of *Yunyangosaurus*.

The neural spine of the axis is posterodorsally oriented (Fig. 3F) as in many other theropods, including *Yangchuanosaurus hepingensis* (ZDM 0024), *Allosaurus*²⁰, and *Sinraptor*¹³. The anterior margin of the neural spine does not extend anteriorly beyond the prezygapophysis, unlike in other non-tetanuran theropods such as *Coelophysis*²⁷, *Dilophosaurus*²⁵, *Carnotaurus*²³, and *Ceratosaurus*²⁴. The anterior margin of the neural spine is thickened toward its anteroventral end, and develops into a ball-like structure. This feature has not been reported in other theropods and may be an autapomorphy of *Yunyangosaurus*.

The preserved anterior postaxial cervical vertebra is platycoelous, with an essentially a flat or weakly convex anterior surface. This also occurs in most non-tetanuran theropods²⁸, the piatnitzkysaurids *Condorraptor* and *Piatnitzkysaurus*^{9,29} and the Chinese Jurassic tetanurans '*Szechuanosaurus*' *zigongensis*¹⁷, *Xuanhanosaurus*¹⁸ and figured cervical vertebra of *Chuandongocoelurus* and *Kaijiangosaurus*¹¹. In contrast, most other tetanurans have convex anterior surfaces of their cervical centra²⁸, including *Monolophosaurus*¹⁹, megalosaurids such as *Leshansaurus*¹⁶ and metriacanthosaurids such as *Sinraptor*¹³. The posterior articular surface is concave. The perimeter of the anterior articular surface of the centrum forms a distinct flattened rim (Fig. 4). This condition is widespread among megalosauroids and is a proposed synapomorphy of the group megalosaurids, piatnitzkysaurids and spinosaurids^{10,22,26,30}. The centrum is considerably longer anteroposteriorly than dorsoventrally in lateral view, with a length:height ratio of 1.44. It is slightly wider transversely than dorsoventrally in posterior view (Table 1). The centrum is invaded by two pneumatic foramina on each lateral side, an anterior and a posterior foramen. The anterior foramen is deep, penetrating into the body of the centrum. However, the posterior one is shallow in the anterior cervical vertebra, taking the form of a fossa that does not penetrate the centrum. Nevertheless, a deep, penetrating posterior pneumatic foramen is present in some of the other cervical centra (described below). The posterior pneumatic foramen is widely-distributed among non-tetanuran theropods, but is absent in most tetanurans^{28,31}. The posterior pneumatic foramen is variably present in some individual cervical and pectoral vertebrae of *Piatnitzkysaurus* (PVL 4073). A well-defined sharp-edged fossa of corresponding size and position to the posterior pneumatic foramen is present in a middle cervical vertebrae of *Condorraptor*²⁹. The pneumatic foramina of the anterior cervical centrum of *Yunyangosaurus* are situated at the upper part of the centrum in lateral view, while that are situated at the middle part of the subsequent centra. The ventral surface of the centrum is mediolaterally thin in ventral view and concave in side view, and bear a low, anteroposteriorly oriented ridge.

The neural arch is invaded ventrolaterally by pneumatic fossae as in many theropods, including *Ceratosaurus*²⁴, *Torvosaurus*^{30,32}, *Piatnitzkysaurus*⁹, *Allosaurus*²⁴, *Carnotaurus*²³, and *Sinraptor*¹³. In *Allosaurus*²⁰ and *Monolophosaurus*¹⁹, the lateral fossae of anterior cervical neural arches are shallow compared to those of *Sinraptor*¹³ and *Yunyangosaurus puanensis*. A horizontal interprezygapophyseal lamina connects the prezygapophyses on the anterior surface of the neural arch. This lamina forms the dorsal roof of the neural canal anteriorly, and the ventral floor of a triangular fossa on the anteroventral surface of the neural arch, together with the




Figure 5. Three middle cervical vertebrae of *Yunyangosaurus puanensis* holotype. The anterior one (PV3, Table 1) in anterior (A), left lateral (B), in posterior (C) and right lateral (D) views; the middle one (PV4, Table 1) in dorsal view (E); the posterior one (PV5, Table 1) in anterior (F), left lateral (G), posterior (H), right lateral (I) and dorsal (J) views. cpol centropostzygapophyseal lamina; dp, diapophysis; ep, epipophysis; ipol, intrapostzygapophyseal lamina; nc, neural canal; ns, neural spine; pl, pneumatic foramen (pleurocoel); poz, postzygapophysis; pp, parapophysis; prdl, prezygodipophyseal lamina; prz, prezygapophysis; spol, spinopostzygapophyseal lamina; sprl, spinoprezygapophyseal lamina. Scale bar = 5 cm.

spinoprezygapophyseal laminae. The spinopostzygapophyseal laminae and the interpostzygapophyseal laminae bound a much broader, rhomb-shaped fossa on the posterior surface of the neural spine.

The diapophyses are located at the fusion line of the centrum and neural arch. They are inclined ventrolaterally as in other theropods such as *Sinraptor* and *Piatnitzkysaurus*^{9,13}. The articular surface of prezygapophysis is approximately circular, while the articular surface of postzygapophysis is a long oval shape. The epipophyses are prominent, extending well beyond the posterior margins of the postzygapophyses, and they are oriented posteriorly rather than posterolaterally as in most other theropods^{9,19,20,30,33}. The presence of prominent and strongly posteriorly-directed epipophyses of the anterior cervical vertebrae may therefore be an autapomorphy of *Yunyangosaurus*. The neural spine is plate-like and posterodorsally oriented, proportionally shorter than in *Ceratosaurus*²⁴ and *Allosaurus*²⁰. The prespinal and postspinal ridges are prominent.

The middle cervical vertebrae are platycoelous (Fig. 5). The centra are slightly longer anteroposteriorly than dorsoventrally in lateral view (Table 1). The posterior articular surfaces are wider transversely than high dorsoventrally, whereas the anterior articular surfaces are smaller, and are higher dorsoventrally than the posterior articular surfaces (Fig. 5B,F) (Table 1). As with the anterior cervicals, the centra are invaded by two pneumatic foramina on each side. But unlike the anterior cervicals, the ventral surfaces of middle cervical vertebrae are broad and flat. The neural arch of the third cervical vertebra is fairly complete, and the upper margin of the neural canal connects the prezygapophyses forming an elongated fossa with the spinoprezygapophyseal laminae in anterior view. The prezygapophysis of the cervical cm3 are large and its articular surfaces face anterodorsolaterally. The postzygapophyses of the cervical cm3 are large with well-developed quadrangular articular surfaces, and prominent epipophyses overhang the postzygapophyses. The neural spines are club-shaped and vertical. They are relatively slender mediolaterally and anteroposteriorly, resulting in a finger-like appearance.

The posterior cervicals are also platycoelous (Figs. 6 and 7). Paired fossae are located ventrally on the anterior surface of the neural spine. These fossae are bounded by the spinoprezygapophyseal laminae laterally and the prezygapophyses ventrally, and are separated on the midline by the prespinal ridge of the neural spine (Fig. 7A). Paired fossae are also present on the posterior surface of the neural arch, bounded ventrally by the dorsal margin of the neural canal, the centropostzygapophyseal laminae and the intrapostzygapophyseal laminae. These fossae are deeper in the penultimate cervical than in the posteriormost one (Figs. 6C and 7C). The centra of the posterior cervicals are invaded by only one pneumatic foramen (the anterior pneumatic foramen) on each lateral side. The posterior pneumatic foramen is absent, unlike in more anterior cervicals. The pneumatic foramina are entirely circular and are deep, penetrating well into the body of the centrum and invading the posterior surfaces of the parapophyses. The articular surfaces of prezygapophyses are approximately circular and dorsolateral, while the articular surfaces of postzygapophyses are long oval and ventrolateral. The epipophyses are extend posteriorly to a level that is slightly posterior to the postzygapophyses. The prespinal and postspinal ridge are more prominent in the posteriormost cervical. The diapophysis originates just dorsal to the level of the neural canal and curves ventrolaterally along its length (Fig. 7F). Three large fossae are located on the lateral surface of the neural arch, ventral to the base of the diapophysis: the infraprezygapophyseal fossa, infradiapophyseal fossa and intrapostzygapophyseal fossa. These fossae are bounded by the prezygodipophyseal lamina, the paraprezygapophyseal lamina, the paradiapophyseal lamina, the posterior centrodipophyseal lamina, the intrapostzygapophyseal lamina and the postzygadiapophyseal lamina. The infraprezygapophyseal fossa and intrapostzygapophyseal fossa are




Figure 6. Penultimate cervical vertebra (PV6, Table 1) of *Yunyangosaurus puanensis* holotype in anterior (A), left lateral (B), posterior (C), right lateral (D), dorsal (E), ventral (F) and 3D image (G) views. acdl, anterior centrodiapophyseal lamina; cpol, centropostzygapophyseal lamina; cpri, centroprezygapophyseal lamina; ep, epiphysis; f, fossa; ho, hyosphene; ipol, intrapostzygapophyseal lamina; nc, neural canal; ns, neural spine; pcdl, posterior centrodiapophyseal lamina; pl, pneumatic foramen (pleurocoel); poz, postzygapophysis; pp, parapophysis; prdl, prezygodipophyseal lamina; prz, prezygapophysis. The red arrow directs bifurcation of the neural spine. The red arrow directs bifurcation of the neural spine. Scale bar = 5 cm.

irregularly quadrilateral, but the infradiapophyseal fossa is triangular (Fig. 7D). In anterior view, the bottom of the right postzygapophyses lamina develop two large and deep fossae divided by a prominent ridge, while only one small and deep fossa developed beneath the bottom of the left postzygapophyses lamina. In the posterior view, the bottom of each postzygadiapophyseal lamina develop a deep fossa, and an obvious ridge can be seen beneath the bottom of the right postzygadiapophyseal lamina (Fig. 7A,C). The neural spines of the posterior cervicals are slender in lateral view and their apices are slightly bifurcate (Figs. 6C, 7A,C).

The three anteriormost dorsal vertebrae of *Yunyangosaurus* have platycoelous centra. This is similar to the condition in most theropods, although the first four or five dorsals of *Acrocanthosaurus* may be opisthocelous³⁴. The ratio of the anteroposterior length of centrum to the height of the posterior articular surface of centrum is 0.85 to 1.13 among anterior dorsal vertebrae of *Yunyangosaurus*, and the ratio of height to width of the posterior articular surface of centrum is 0.85 to 0.92. The anterior articular surfaces of the centra are marked by a well-developed circumferential groove, and two shallow fossae are also present on the anterior articular surfaces of all dorsals, located just dorsal to mid-height (Figs. 8D,J and 9A). The anterior articular surfaces of the dorsals are roughly circular and posterior articular surfaces are nearly square.

The central portion of each dorsal centrum is transversely constricted relative to the articular surfaces. Distinct, single pneumatic foramina invade the lateral surfaces of the centra posterior to the parapophyses in anterior dorsals. Pneumatic foramina are present only in the cervical and anterior dorsal centra of *Yunyangosaurus*, and are absent from more posterior dorsals. This is similar to the condition in many Jurassic theropods³⁵, for example, the posteriormost centrum to bear a pneumatic foramen is the 14th presacral of *Sinraptor*¹³ and the 13th presacral of *Allosaurus*²⁰. Nevertheless, pneumatic foramina in more posterior dorsal centra are present in geologically younger members of many groups of theropods and take various morphologies³⁵. For example, large pneumatic openings are found in all dorsal centra of *Torvosaurus*³⁰. There is one conspicuous fossa beneath each parapophysis in all of the preserved dorsals of *Yunyangosaurus*. Radial grooves are present on the posteroventral surface of the centrum in anterior dorsals (Fig. 8H,I). The hypapophyses are also prominent in the anterior dorsals. Prominent hypoapophyses are widely-distributed among adult tetanurans, including *Allosaurus*²⁰, *Condoraptor*²⁹, *Marshosaurus* (CMNH 21704), *Sinraptor*¹³, *Piatnitzkysaurus*⁹ and *Streptospondylus*³⁶. The most prominent hypapophysis of *Allosaurus* is in the second dorsal (Madsen, 1976), which is the 11th presacral vertebra, as in *Sinraptor*¹³.




Figure 7. Last cervical vertebra (PV7, Table 1) of *Yonyangosaurus puanensis* holotype in anterior (A), left lateral (B), posterior (C), right lateral (D), dorsal (E) and ventral (F). acdl, anterior centrodiapophyseal lamina; cpol centropostzygapophyseal lamina; cprl, centroprezygapophyseal lamina; dp, diapophysis; ep, epiphysis; idpf, infradiapophyseal fossa; ipol, intrapostzygapophyseal lamina; ipozf, infrapostzygapophyseal fossa; iprzf, infraprezygapophyseal fossa; nc, neural canal; ns, neural spine; pcdl, posterior centrodiapophyseal lamina; pl, pneumatic foramen (pleurocoel); podl, postzygadiapophyseal lamina; posr, postspinal ridge; poz, postzygapophysis; pp, parapophysis; prdl, prezygodipophyseal lamina; prsr, prespinal lamina; prz, prezygapophysis; spol, spinopostzygapophyseal lamina; sprl, spinoprezygapophyseal lamina. The red arrows direct bifurcation of the neural spine. Scale bar = 5 cm.

The hyposphenes are large in the anterior dorsals and have a roughly constant transverse width dorsoventrally. This differs from the ventrally-wide, triangular hyposphene morphology that is found among many early-diverging tetanurans³⁷. Step-like ridges are present lateral to the hyposphenes, running posterodorsally from the dorsal border of neural canal to the posterior edge of postzygapophyses (Figs. 6C and 8C). These ridges have a homoplastic distribution among early-diverging tetanurans being present both among megalosauroids and in metriacanthosaurids³⁷. The spinopostzygapophyseal lamina extends posterodorsally to the posterolateral margin of the neural spine from the posterior corner of the postzygapophysis. The prespinal and postspinal ridges are developed in the anterior dorsals (Fig. 8A,C). The spinopostzygapophyseal laminae are developed in the anterior dorsals. The articular surface of the diapophysis is trapeziform and slightly concave (Fig. 8B). Below the base of the diapophysis, there are three large fossae called infraprezygapophyseal fossa, infradiapophyseal fossa and infrapostzygapophyseal fossa, which on the lateral surface of the base of the neural arch. These fossae are formed by the prezygodipophyseal lamina, the paraprezygapophyseal lamina, the paradiapophyseal lamina, the posterior centrodiapophyseal lamina, the intrapostzygapophyseal lamina and the postzygadiapophyseal lamina. The infraprezygapophyseal fossa and infrapostzygapophyseal fossa are irregularly quadrilateral, but the infradiapophyseal fossa is triangular (Fig. 8B). The bottom of the prezygodipophyseal laminae develop two small ridges (one is the prezygodiapophyseal ridge) and some small shallow fossae in the anterior dorsal. The neural spine is claviform in the anterior dorsals and the tops of the anterior dorsals are transversely bifurcated as in the posterior cervicals, but the bifurcation is deeper than in the latter. The articular surface of prezygapophyses are approximately circular, while the articular surface of postzygapophyses are oval.

The centrum of the preserved anterior middle dorsal is amphiplatyan. The amphiplatyan condition is present in all dorsals of *Carnotaurus*²³, whereas it is only present from the fifth dorsal (14th presacral) and more posteriorly in *Allosaurus*²⁰ and from the sixth (15th presacral) of *Monolophosaurus*¹⁹. Clear, radial grooves can be found in the anteroventral and posteroventral portions of the centrum (Fig. 9F). A shallow depression is present on the lateral surface of this centrum in *Yonyangosaurus*. This may be homologous to the large, extensive, fossa-like pneumatic foramina that are present in the anterior dorsal centra of some megalosauroids such as *Torvosaurus* and *Eustreptospondylus*^{10,22,30}, though it is relatively shallow, and not pronounced as in those taxa.

The neural canal of the anterior middle dorsal is 11 mm high and 8 mm wide while the anterior articular surface is 23 mm high and 25 mm wide, so the neural canal is about 1/3 to 1/2 as big as the centrum (Fig. 9A). The hypantrum is highly developed and the ridges of the middle portion between the dorsal margin of the neural canal and the junction of the prezygapophyses are developed. These ridges are called intraprezygapophyseal laminae. They contact the articular surfaces of the prezygapophyses, forming two triangular fossae with the margin




Figure 8. Three anteriormost dorsal vertebrae of *Yunyangosaurus puanensis* holotype. Dorsal 1 (PV9, Table 1) in anterior (A), left lateral (B), in posterior (C); Dorsal 2 (PV8, Table 1) in anterior (D), right lateral (E), ventral (F); Dorsal 3 (PV10, Table 1) in left lateral (G), ventral (H); Dorsal 3 (PV10, Table 1) in ventral (I), anterior (J), right lateral (K), posterior (L) and 3D images (M–O). acdl, anterior centrodiapophyseal lamina; cpol centropostzygapophyseal lamina; cprl, centroprezygapophyseal lamina; dp, diapophysis; f, fossa; hy, hypapophysis; ho, hyposphene; idpf, infradiapophyseal fossa; ipozf, infrapostzygapophyseal lamina; iprpf, infraprezygapophyseal lamina; iprl, infraprezygapophyseal lamina; nc, neural canal; ns, neural spine; pcdl, posterior centrodiapophyseal lamina; pl, pneumatic foramen (pleurocoel); posr, postspinal ridge; poz, postzygapophysis; pp, parapophysis; prdl, prezygodipophyseal lamina; prsr, prezygodipophyseal lamina; prz, prezygapophysis; spol, spinopostzygapophyseal lamina. The red arrows direct bifurcation of the neural spine. Scale bar = 5 cm.




Figure 9. An anterior middle dorsal vertebra (PV11, Table 1) of *Yunyangosaurus puanensis* holotype in anterior (A), left lateral (B), posterior (C), right lateral (D), dorsal (E), ventral (F) and 3D image (G). acdl, anterior centrodiapophyseal lamina; cprl, centroprezygapophyseal lamina; dp, diapophysis; f, fossa; idpf, infradiapophyseal fossa; ipol, intrapostzygapophyseal lamina; ipozf, infrapostzygapophyseal fossa; iprpf, infraprezygapophyseal fossa; nc, neural canal; ns, neural spine; pcdl, posterior centrodiapophyseal lamina; pl, lateral depression (pleurocoel); pp, parapophysis; prdl, prezygodipophyseal lamina; prdr, prezygodipophyseal ridge; prz, prezygapophysis; spd, spinodiapophyseal lamina; spol, spinopostzygapophyseal lamina; spor, spino-spinopostzygapophyseal ridge; sprl, spinoprezygapophyseal lamina. Scale bar = 5 cm.

of the neural canal and anterior centroprezygapophyseal laminae (Fig. 9A). The hyposphene is smaller in the anterior middle dorsal than in the anterior dorsals. A ridge is visible on the spinopostzygapophyseal lamina of the anterior/middle dorsal, while it is missing in the anterior dorsal (Fig. 9C). The spinoprezygapophyseal and spinopostzygapophyseal laminae are developed in the anterior middle dorsal. The spinoprezygapophyseal lamina is obvious in the anterior middle dorsal but only a gentle ridge in the anterior dorsals (Fig. 9A). The diapophyses are upswept and below the base of which are three large fossae called infraprezygapophyseal fossa, infradiapophyseal fossa and infrapostzygapophyseal fossa, on the lateral surface of the base of the neural arch. These fossae are formed by the prezygodipophyseal lamina, the paraprezygapophyseal lamina, the paradiapophyseal lamina, the posterior centrodiapophyseal lamina, the intrapostzygapophyseal lamina and the postzygodipophyseal lamina. The infraprezygapophyseal fossa and infrapostzygapophyseal fossa are irregularly quadrilateral, but the infradiapophyseal fossa is triangular (Fig. 9D). In the infrapostzygapophyseal fossa, the centropostzygapophyseal ridge is visible in the anterior middle dorsal but illegible in the anterior dorsals (Fig. 9B). The neural spine is incompletely preserved, but it is inferred to be plate-like (Fig. 9A).

Two proximal portions of the cervical ribs are preserved. The capitulum and tuberculum are widely separated, and the articular facets lie in the same plane, so presumably they belong to posterior cervicals. Deep fossae are found between the base of the capitulum, tuberculum, and prominent anterolateral process. The angle between the capitulum and tuberculum is about 70°. Some partial dorsal ribs are preserved. The capitulum and tuberculum are more widely separated, and the proximal pneumatopore is absent. The cross-sectional shapes of the ribs are subcircular.

Three unforked chevrons with bridged haemal canal are preserved, and the haemal canal is short approximately 20% chevron length (Fig. 10).

Discussion

Yunyangosaurus puanensis is represented by limited material, but it displays numerous informative features for its systematic position and taxonomy. It is highly likely a tetanuran dinosaur based on the general morphology of the presacral vertebrae, and in particular several tetanuran or orinoidan synapomorphies are present^{10,38}: the axial neural spine is narrow anteroposteriorly and somewhat rod-like; the prezygapophyses of anterior cervical vertebrae are enlarged and situated entirely lateral to the neural canal; and there is a pronounced ventral keel in the anterior dorsal vertebrae.

However, *Yunyangosaurus puanensis* also lacks several derived features seen in most other early-branching tetanurans^{10,38}. These features include strongly opisthocoelous cervical centra, cervical vertebrae with only one pair of pneumatic foramina (rather than two, including both the anterior and posterior pneumatic foramen, as in




Figure 10. A partial chevron (A1-A5) and two ribs (B1-B2 are dorsal ribs, C1-C3 are cervical ribs) of *Yunyangosaurus puanensis* holotype.

most non-tetanuran theropods), and the presence of prominent axial spinopostzygapophyseal laminae. Although the anterior surfaces of the anterior cervical centra of *Yunyangosaurus puanensis* are weakly convex, this is unlike the prominent, convexity seen in the opisthocoelous condition that is widespread among early-diverging tetanurans. Furthermore, the middle and posterior cervical vertebrae of *Yunyangosaurus* are strictly platycoelous with flat anterior surfaces. Among tetanurans, this morphology, with flat to weakly convex anterior surfaces of the cervical centra, is also present in ‘*Szechuanosaurus*’ *zigongensis* (ZDM 9011), the piatnitzkysaurid *Piatnitzkysaurus*⁹, *Xuanhanosaurus*¹⁸ and *Condorraptor*³⁹. Furthermore, the presence of a second, posterior, pneumatic foramen is variably present in some cervical centra of *Piatnitzkysaurus*³⁵, although it is much more prominently and consistently developed in *Yunyangosaurus*, similar to the condition in most non-tetanuran theropods²⁸. The axial spinopostzygapophyseal lamina is reduced in most tetanurans, unlike in *Yunyangosaurus*. Among tetanurans, a prominent axial spinopostzygapophyseal lamina is also present in metriacanthosaurids such as *Sinraptor*¹³.

Yunyangosaurus puanensis also lacks an axial intercentrum with an anterodorsally inclined ventral surface. This feature is present in many early-branching tetanurans including *Monolophosaurus*¹⁹ and *Sinraptor*¹³. However, the ventral surface of the axial intercentrum of *Yunyangosaurus* is horizontal similar to the condition in some megalosauroids, such as *Leshansaurus* and *Piatnitzkysaurus*^{9,16}.

Yunyangosaurus possesses several features that are only known in some megalosauroids, especially piatnitzkysaurids (described above). These also include the presence of a prominent rim round articular surface of cervicals¹⁰ and bifurcated anterior dorsal neural spines (present in piatnitzkysaurids)⁷. A numerical cladistic analysis was then ran to assess this phylogenetic hypothesis (see Methods), but which fails to place *Yunyangosaurus* within the Megalosauroida and instead posits it in an unresolved position with many other tetanurans (Fig. 11). Noteworthy is also that the inclusion of *Yunyangosaurus* in the dataset led to collapsing of the Megalosauroida and the Coelurosauria. This suggests the complex combination of character states in *Yunyangosaurus* has an impact on the tetanuran phylogeny, which is still far from robust for understanding the evolution of the group. However, we reran the analysis using implied weighting (see Methods), and this analysis places *Yunyangosaurus* within the Megalosauroida in an unresolved position with many other megalosauroids (Fig. 12).

The discovery of *Yunyangosaurus* has implication for our understanding of character evolution of tetanuran theropods. For example, *Yunyangosaurus* displays several features that are generally absent in most other tetanurans, and these features include the presence of posterior pneumatic foramina throughout the cervical series and in particular pneumatic foramen in middle cervical centra (also present occasionally in some cervical centra of piatnitzkysaurids), prominent spinopostzygapophyseal laminae (also present in metriacanthosaurids), and flat anterior articular surfaces of the cervical centra (also present in piatnitzkysaurids).

However, currently our knowledge of *Yunyangosaurus* is relatively incomplete, preventing us from attaining a more complete understanding of its affinities among early tetanurans. For now, *Yunyangosaurus* serves to highlight the potentially complex character distribution of character states of the axial column among early-branching tetanurans, and especially among Chinese Jurassic tetanurans.




Figure 11. Strict consensus of 4860 most parsimonious trees produced by our analysis showing the systematic position of *Yunyangosaurus puanensis*.

Methods

Phylogenetic analysis. We added *Yunyangosaurus* into a comprehensive dataset for tetanuran phylogeny⁴⁰, and the final matrix includes 351 characters and 63 operational taxonomical units (OTUs). The matrix was analyzed using TNT 1.5⁴¹, with the ‘New Technology’ search option, using a driven search that stabilized consensus twice with a factor of 25, and default settings for sectorial, ratchet, tree drift and tree fusion. The resulting 142 most parsimonious trees (MPTs) with a tree length = 1103 step, a CI = 0.39, and a RI = 0.67 were then subjected to tree bisection and reconnection (TBR) branch swapping, and finally 4860 MPTs were found. The strict




Figure 12. Strict consensus of 1107 most parsimonious trees produced by our parsimonious analysis with implied weighting showing the systematic position of *Yonyangosaurus puanensis*.

consensus of these 4860 MPTs places *Yonyangosaurus* at the base of the Tetanurae in an unsolved position with several other early-branching tetanuran theropods. However, we reran the analysis using implied weighting parsimony with a k value of 8.0, and this analysis places *Yonyangosaurus* within the Megalosauroidea in an unresolved position with many other megalosauroids (Fig. 12).

Received: 24 June 2019; Accepted: 25 November 2019;
 Published online: 10 January 2020

References

- Dong, Z. M., Zhou, S. & Zhang, Y. The dinosaurian ramians from Sichuan Basin. *China. Paleontol Sinica, n.s. C* **23**, 1–145 (1983).
- Peng, G., Ye, Y., Gao, Y., Shu, C. & Jiang, S. *Jurassic dinosaur faunas in Zigong*. (Sichuan Renmin Press, Chengdu, 2005).
- Gao, Y. *Yangchuanosaurus hepingensis* - a new species of carnosaur from Zigong, Sichuan. *Vertebrata Palasiatica* **30**, 313–324 (1992).
- Dong, Z. & Tang, Z. A new Mid-Jurassic theropod (*Gasosaurus constructus* gen. et sp. nov.) from Dashanpu, Zigong, Sichuan Province, China. *Vertebrata Palasiatica* **23**, 77–83 (1985).
- Huang, D. Jurassic integrative stratigraphy and timescale of China. *Science China Earth Sciences* **62**, 223–255 (2019).
- Fang, Y., Liu, Y., Zhang, Y. & Wang, B. Elemental analysis and calcification characteristics of ostracods from the Middle Jurassic Xintiangou Formation in eastern Sichuan province, China (in Chinese). *Acta Palaeontologica Sinica* **51**, 96–101 (2012).
- Wang, S. *et al.* Extreme ontogenetic changes in a ceratosaurian theropod. *Current Biology* **27**, 144–148 (2017).
- Currie, P. & Xijin, Z. A new carnosaur from the Jurassic of Xinjiang, People's Republic of China. *Canadian Journal of Earth Sciences* (1993).
- Bonaparte, J. F. Les dinosaures (carnosaures, allosauridés, sauropodes, cétiosaures) du Jurassique Moyen de Cerro Cóndor (Chubut, Argentine). *Annales de Paléontologie* **72**, 247–289 (1986).
- Carrano, M. T., Benson, R. B. & Sampson, S. D. The phylogeny of Tetanurae (Dinosauria: Theropoda). *Journal of Systematic Palaeontology* **10**, 211–300 (2012).
- He, X. *The Vertebrate Fossils of Sichuan (in Chinese)*. (Sichuan Science and Technology Press, 1984).
- Gao, Y. A complete carnosaur skeleton from Zigong, Sichuan: *Yangchuanosaurus hepingensis*. *Sichuan Science and Technology Press, Chengdu*, 1–100 (1999).
- Currie, P. J. & Zhao, X. A new carnosaur (Dinosauria, Theropoda) from the Jurassic of Xinjiang, People's Republic of China. *Canadian Journal of Earth Sciences* **30**, 2037–2081 (1993).
- Dong, Z., Zhang, Y., Li, X. & Zhou, S. A new carnivorous dinosaur of Yongchuan, Sichuan. *Chinese Science Bulletin* **23**, 302–304 (1978).
- Wu, X., Currie, P. J., Dong, Z., Pan, S. & Wang, T. A new theropod dinosaur from the Middle Jurassic of Lufeng, Yunnan, China. *Acta Geologica Sinica-English Edition* **83**, 9–24 (2009).
- Li, F., Peng, G., Ye, Y., Jiang, S. & Huang, D. A new carnosaur from the Late Jurassic of Qianwei, Sichuan, China. *Acta Geologica Sinica* **83**, 1203–1213 (2009).
- Gao, Y. A new species of *Szechuanosaurus* from the Middle Jurassic of Dashanpu, Zigong, Sichuan. *Certebrata Palasiatica*, **6** (1993).
- Dong, Z., Zhou, S. & Zhang, Y. Dinosaurs from the Jurassic of Sichuan basin, China. *Palaeontologia Sinica (new series C)* **162**, 1–145 (1983).
- Zhao, X., Benson, R. B., Brusatte, S. L. & Currie, P. J. The postcranial skeleton of *Monolophosaurus jiangi* (Dinosauria: Theropoda) from the Middle Jurassic of Xinjiang, China, and a review of Middle Jurassic Chinese theropods. *Geological Magazine* **147**, 13–27 (2009).
- Madsen, J. H. Jr. *Allosaurus fragilis*: a revised osteology. *Utah Geological and Mining Survey Bulletin* **109**, 1–163 (1976).
- Harris, J. D. A Reanalysis of *Acrocanthosaurus atokensis*, its Phylogenetic Status, and Paleobiogeographic Implications, Based on a New Specimen from Texas: *Bulletin 13*. Vol. 13 (New Mexico Museum of Natural History and Science, 1998).
- Sadleir, R., Barrett, P. M. & P. P. H. The anatomy and systematics of *Eustreptospondylus oxoniensis*, a theropod dinosaur from the Middle Jurassic of Oxfordshire, England. *Monograph of the Palaeontographical Society, London* **162**, 1–82 (2008).
- Bonaparte, J. F., Novas, F. E. & Coria, R. A. *Carnotaurus sastrei* Bonaparte, the horned, lightly built carnosaur from the Middle Cretaceous of Patagonia. *Contrib. in sci./Natural history museum of Los Angeles county* **419**, 1–41 (1990).
- Gilmore, C. W. *Osteology of the carnivorous Dinosauria in the United States National museum: with special reference to the genera Antrodemus (Allosaurus) and Ceratosaurus*. (US Government printing office, 1920).
- Welles, S. P. *Dilophosaurus wetherilli* (Dinosauria, Theropoda). Osteology and comparisons. *Palaeontographica Abteilung A*, 85–180 (1984).
- Charig, A. J. & Milner, A. C. *Baryonyx walkeri*, a fish-eating dinosaur from the Wealden of Surrey. *Bulletin-Natural History Museum Geology Series* **53**, 11–70 (1997).
- Colbert, E. H. *The triassic dinosaur Coelophysis*. (Museum of Northern Arizona, 1989).
- Rauhut, O. W. M. The interrelationships and evolution of basal theropods (Dinosauria, Saurischia). *Palaeontology* **69**, 1–215 (2003).
- Rauhut, O. W. Osteology and relationships of a new theropod dinosaur from the Middle Jurassic of Patagonia. *Palaeontology* **48**, 87–110 (2005).
- Britt, B. Theropods of Dry Mesa Quarry (Morrison Formation, late Jurassic), Colorado, with emphasis on the osteology of *Torvosaurus tanneri*. *Brigham Young University Geology. Studies* **37**, 1–72 (1991).
- Gauthier, J. Saurischian monophyly and the origin of birds. *Memoirs of the California Academy of sciences* **8**, 1–55 (1986).
- Britt, B. B. *Pneumatic postcranial bones in dinosaurs and other archosaurs*, University of Calgary, Alberta, (1993).
- Madsen, J. Jr. & Welles, S. *Ceratosaurus* (Dinosauria, Theropoda). A revised osteology. *Utah Geological Survey. Miscellaneous Publications* **2**, 1–80 (2000).
- Stovall, J. W. & Langston, W. *Acrocanthosaurus atokensis*, a new genus and species of Lower Cretaceous Theropoda from Oklahoma. *The American Midland Naturalist* **43**, 696–728 (1950).
- Benson, R. B. J., Butler, R. J., Carrano, M. T. & O'Connor, P. M. Air-filled postcranial bones in theropod dinosaurs: physiological implications and the 'reptile'-bird transition. *Biological Reviews* **87**, 168–193 (2012).
- Allain, R. R. de *Streptospondylus altdorfensis*, le dinosaure théropode de Cuvier, du Jurassique de Normandie. *Geodiversitas* **23**, 349–367 (2001).
- Benson, R. B. J. A description of *Megalosaurus bucklandii* (Dinosauria: Theropoda) from the Bathonian of the UK and the relationships of Middle Jurassic theropods. *Zoological Journal of the Linnean Society* **158**, 882–935 (2010).
- Rauhut, O. W. M. The interrelationships and evolution of basal theropod dinosaurs. *Palaeontology* **69**, 1–215 (2003).
- Rauhut, O. W. & Xu, X. The small theropod dinosaurs *Tugulusaurus* and *Phaedrolosaurus* from the Early Cretaceous of Xinjiang, China. *Journal of Vertebrate Paleontology* **25**, 107–118 (2005).
- Rauhut, O. W., Hübner, T. & Lanser, K.-P. A new megalosaurid theropod dinosaur from the late Middle Jurassic (Callovian) of north-western Germany: Implications for theropod evolution and faunal turnover in the Jurassic. *Palaeontologia Electronica* **19**, 1–65 (2016).
- Goloboff, P. A., Farris, J. S. & Nixon, K. C. TNT, a free program for phylogenetic analysis. *Cladistics* **24**, 774–786 (2008).

Acknowledgements

We thank Wei Guangbiao and He Jianjun for coordinating the fieldwork in Yunyang, Han Xiangsong for collecting the specimen, Zhang Yuqing for preparing the specimen. This work is supported by Natural Science Foundation of Chongqing, China (No. cstc2018jcyjAX0435), Postdoctoral Project of Chongqing, China (No. Xm2017069) and Project of Dinosaur Fossil Protection and Research of Chongqing Planning and Natural Resources Bureau (No. kj-2018035). X.X is supported by the National Natural Science Foundation of China (Grant No. 41688103).

Author contributions

Hui Dai, Xing Xu and Roger Benson wrote the main manuscript text. Qingyu Ma, Shan Jiang, Deliang Li and Guangzhao Peng revised and improved the manuscript. Hui Dai, Qingyu Ma, Haiqian Hu and Yuxuan Zhou prepared all figures. Xufeng Hu, Chao Tan, Ming Xiao, Feng Zhang and Yilun Yu oversaw the project. Ning Li and Zhaoying Wei assisted with the measurement of the specimens and the writing of the manuscript. Yilun Yu helped to revise the phylogenetic tree. All authors reviewed the manuscript.

Competing interests

The authors declare no competing interests.

Additional information

Supplementary information is available for this paper at <https://doi.org/10.1038/s41598-019-56959-x>.

Correspondence and requests for materials should be addressed to H.D. or X.X.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2020

Terms and Conditions

Springer Nature journal content, brought to you courtesy of Springer Nature Customer Service Center GmbH (“Springer Nature”).

Springer Nature supports a reasonable amount of sharing of research papers by authors, subscribers and authorised users (“Users”), for small-scale personal, non-commercial use provided that all copyright, trade and service marks and other proprietary notices are maintained. By accessing, sharing, receiving or otherwise using the Springer Nature journal content you agree to these terms of use (“Terms”). For these purposes, Springer Nature considers academic use (by researchers and students) to be non-commercial.

These Terms are supplementary and will apply in addition to any applicable website terms and conditions, a relevant site licence or a personal subscription. These Terms will prevail over any conflict or ambiguity with regards to the relevant terms, a site licence or a personal subscription (to the extent of the conflict or ambiguity only). For Creative Commons-licensed articles, the terms of the Creative Commons license used will apply.

We collect and use personal data to provide access to the Springer Nature journal content. We may also use these personal data internally within ResearchGate and Springer Nature and as agreed share it, in an anonymised way, for purposes of tracking, analysis and reporting. We will not otherwise disclose your personal data outside the ResearchGate or the Springer Nature group of companies unless we have your permission as detailed in the Privacy Policy.

While Users may use the Springer Nature journal content for small scale, personal non-commercial use, it is important to note that Users may not:

1. use such content for the purpose of providing other users with access on a regular or large scale basis or as a means to circumvent access control;
2. use such content where to do so would be considered a criminal or statutory offence in any jurisdiction, or gives rise to civil liability, or is otherwise unlawful;
3. falsely or misleadingly imply or suggest endorsement, approval, sponsorship, or association unless explicitly agreed to by Springer Nature in writing;
4. use bots or other automated methods to access the content or redirect messages
5. override any security feature or exclusionary protocol; or
6. share the content in order to create substitute for Springer Nature products or services or a systematic database of Springer Nature journal content.

In line with the restriction against commercial use, Springer Nature does not permit the creation of a product or service that creates revenue, royalties, rent or income from our content or its inclusion as part of a paid for service or for other commercial gain. Springer Nature journal content cannot be used for inter-library loans and librarians may not upload Springer Nature journal content on a large scale into their, or any other, institutional repository.

These terms of use are reviewed regularly and may be amended at any time. Springer Nature is not obligated to publish any information or content on this website and may remove it or features or functionality at our sole discretion, at any time with or without notice. Springer Nature may revoke this licence to you at any time and remove access to any copies of the Springer Nature journal content which have been saved.

To the fullest extent permitted by law, Springer Nature makes no warranties, representations or guarantees to Users, either express or implied with respect to the Springer nature journal content and all parties disclaim and waive any implied warranties or warranties imposed by law, including merchantability or fitness for any particular purpose.

Please note that these rights do not automatically extend to content, data or other material published by Springer Nature that may be licensed from third parties.

If you would like to use or distribute our Springer Nature journal content to a wider audience or on a regular basis or in any other manner not expressly permitted by these Terms, please contact Springer Nature at

onlineservice@springernature.com