

Short communication

First definitive ankylosaurian dinosaur from the Cretaceous of Jilin Province, northeastern China

Wenjie Zheng^{a, b, *}, Masateru Shibata^{c, d}, Chun-Chi Liao^{e, f, g}, Soki Hattori^{c, d}, Dongchun Jin^h, Changzhu Jin^e, Xing Xu^{e, f}^a Zhejiang Museum of Natural History, Hangzhou, Zhejiang 310014, People's Republic of China^b State Key Laboratory of Palaeobiology and Stratigraphy (Nanjing Institute of Geology and Palaeontology, CAS, People's Republic of China), Nanjing, Jiangsu 210008, People's Republic of China^c Institute of Dinosaur Research, Fukui Prefectural University, Matsuoka Kenjojima, Eiheiji, Fukui 910-1195, Japan^d Fukui Prefectural Dinosaur Museum, Muroko, Terao, Katsuyama, Fukui 911-8601, Japan^e Key Laboratory of Vertebrate Evolution and Human Origins of Chinese Academy of Sciences, Institute of Vertebrate Paleontology and Paleoanthropology, Chinese Academy of Sciences, Beijing 100044, People's Republic of China^f CAS Center for Excellence in Life and Palaeoenvironment, Beijing 100044, People's Republic of China^g University of Chinese Academy of Sciences, Beijing 100049, People's Republic of China^h Yanbian Seismological Bureau, Yanji, Jilin 133001, People's Republic of China

ARTICLE INFO

Article history:

Received 1 February 2021

Received in revised form

7 June 2021

Accepted in revised form 8 July 2021

Available online 14 July 2021

Keywords:

Ankylosauria

Cretaceous

Longjing Formation

Jilin Province

ABSTRACT

We here report an ankylosaurian ilium from the Albian–Cenomanian Longjing Formation of Yanji City, Jilin Province, northeastern China. The diagnostic features allowing referral of this specimen to Ankylosauria include: an ilium rotated horizontally dorsal to the acetabulum with the primitive lateral surface facing ventrally; long and laterally divergent preacetabular process; and a shallow, cup-like acetabulum. This new specimen represents the first definitive ankylosaurian dinosaur from Jilin Province, and the easternmost ankylosaurian occurrence in China. The present discovery demonstrates the potential for future finds in the Longjing Formation, especially in the Longshan fossil beds of Yanji City, such as additional ankylosaurian or other dinosaur remains that might provide significant data on dinosaur evolution during the middle Cretaceous of eastern Asia.

© 2021 Elsevier Ltd. All rights reserved.

1. Introduction

Ankylosauria is a clade of quadrupedal herbivorous armored dinosaurs, which are known from all continents with the exception of Africa (Vickaryous et al., 2004). The Asian ankylosaurian dinosaurs are known mostly from China and Mongolia (Arbour and Currie, 2016). In northeastern China, ankylosaurians have only been reported from Liaoning Province (Amiot et al., 2010; Dong, 2002; Han et al., 2014; Ji et al., 2014; Lü et al., 2007a; Xu et al., 2001; Yang et al., 2017).

Yanji City is located in the eastern part of Jilin Province, northeast to Liaoning Province (Fig. 1). The Tongfosi Formation in the west of Yanji has previously produced footprints of several dinosaurian groups, including theropods, ornithopods, and quadrupedal dinosaurs of uncertain affinities (Matsukawa et al.,

1995; Xing et al., 2017). It was not until 2018 that dinosaur body fossils from Yanji were reported (Jin et al., 2018; Zhang et al., 2018). They were mainly discovered in the Longjing Formation of Longshan Mountain, in the south of Yanji, though one specimen was found from the Chaoyang River (Sun River in Jin et al., 2018), which is in the west of Yanji City.

In September 2017, IVPP, Fukui Prefectural University, and FPDM conducted a joint dinosaur expedition in Yanji, together with members of the Yanji Bureau of Land and Resources and the Yanbian Seismological Bureau. Dongchun Jin discovered one partial left ankylosaurian ilium in the east bank of the Chaoyang River during the expedition. It is the first definitive ankylosaurian dinosaur body fossil from Jilin Province.

1.1. Institutional abbreviations

AMNH, American Museum of Natural History, New York, New York, USA; CMN, Canadian Museum of Nature, Ottawa, Ontario,

* Corresponding author. Zhejiang Museum of Natural History, Hangzhou, Zhejiang 310014, People's Republic of China.

E-mail address: zhengwj@zmnh.com (W. Zheng).




Fig. 1. Locality maps of the fossil locality. Map of Jilin Province showing the location of Yanji City (A), and Map of Yanji City with the fossil locality (marked by a star icon) (B).

Canada; FPDM, Fukui Prefectural Dinosaur Museum, Fukui, Japan; IVPP, Institute of Vertebrate Paleontology and Paleoanthropology, Chinese Academy of Sciences, Beijing, China; NHMUK, Natural History Museum, London, United Kingdom; ZMNH, Zhejiang Museum of Natural History, Hangzhou, China.

2. Geological setting

The Yanji Basin is one of the largest terrestrial Mesozoic basins in the east of Jilin Province, with a total area of 1,500 km². The basin has well-exposed Mesozoic strata, especially Cretaceous ones, which include the Tongfosi, Dalazi, and Longjing formations in ascending order (Bureau of Geology and Mineral Resources of Jilin Province, 1988; Jin et al., 2018). The first body fossil was unearthed from the grey-green sandstone of the Longjing Formation, exposed on the east bank of the Chaoyang River, north of Chaoyangchuan Town (Fig. 1B).

The Longjing Formation, composed of purple-red and pale blue pebbled sandstone, and siltstone with nodular marl, is widely distributed around Chaoyangchuan, Mao'ershan, Longjing, and Jingu in the Yanji Basin. This formation reaches approximately 1,355 m in total thickness and is unconformably covered on the Dalazi Formation (Bureau of Geology and Mineral Resources of Jilin Province, 1988). The geological age of the Longjing Formation is still controversial. Although sometimes considered as the late Early Cretaceous in age (Bureau of Geology and Mineral Resources of Jilin Province, 1988; Sun and Zheng, 2000), it is usually placed in the early Late Cretaceous (Bureau of Geology and Mineral Resources of Jilin Province, 1988; Cui et al., 1994; Jin et al., 2018). Biostratigraphic evidence suggests that the Longjing Formation belongs to the early

Late Cretaceous (Bureau of Geology and Mineral Resources of Jilin Province, 1988; Cui et al., 1994; Jin et al., 2018). Zircon and apatite fission track ages of the underlying Dalazi Formation are 100 Ma and 95 Ma, respectively (Li and Gong, 2008), which indicates that the geological age of the Longjing Formation should be younger, and thus dated to the Late Cretaceous. Jin et al. (2018) placed the Longjing Formation in the Cenomanian, based on these radiometric data and stratigraphic correlation. However, recent SIMS U–Pb zircon dating for the uppermost part of the Dalazi Formation in the Yanji Basin gave an age of 105.14 ± 0.37 Ma (Zhong et al., 2020). Moreover, recent CA-ID-TIMS zircon dating from the fossil beds of Longshan Mountain resulted in an age of 101.039 ± 0.061 Ma (Zhang et al., 2018). Both suggest an older age for the Dalazi and Longjing formations. Ostracod-based biostratigraphic data support an Albian–Cenomanian age for this formation (Zhong et al., 2020). Considering these recent researches, we regard a late Albian–Cenomanian age to be reasonable for the Longjing Formation in this study.

3. Systematic paleontology

Dinosauria Owen 1842
 Ornithischia Seeley 1887
 Thyreophora Nopcsa 1915
 Ankylosauria Osborn 1923

Ankylosauria indet.

Material. IVPP V26052, an incomplete left ilium lacking its anterior and posterior ends (Fig. 2).

Locality and horizon. Chaoyang River, Chaoyangchuan Town, Yanji City, Jilin Province, northeast China. Longjing Formation (upper Albian–Cenomanian) (Bureau of Geology and Mineral Resources of Jilin Province, 1988; Jin et al., 2018; Zhang et al., 2018).

4. Description and comparisons

The ilium is incompletely preserved, missing its anterior and posterior ends (Fig. 2). The preserved anteroposterior length is 64.0 cm and the maximum mediolateral width is 25.5 cm. Despite the absence of all other materials, this specimen can be identified as an ankylosaurian because the ilium is one of the most distinctive elements in an ankylosaurian postcranial skeleton (Coombs, 1979). The ilium has several ankylosaur features that distinguish it from those of other dinosaurs, such as the ilium blade rotating horizontally dorsal to the acetabulum with the primitive lateral surface facing ventrally, long, and laterally divergent preacetabular process, and a shallow cup-like acetabulum (Coombs, 1978; Vickaryous et al., 2004).

The dorsal surface of the ilium is rough, with some irregular protuberances, pits, and longitudinal striations. In dorsal view, the ilium is relatively straight, and the lateral edge is slightly concave. The preacetabular process is long and compressed dorsoventrally, as in other ankylosaurians (Vickaryous et al., 2004). The portion dorsal to the acetabulum is domed, with the lateral portion of the ilium rotated slightly ventrally. In lateral view, the acetabulum is visible. The lateral edge of the preacetabular process is slightly curved ventrally towards anteriorly, as in most ankylosaurians. In contrast, the preacetabular process is curved strongly ventrally in some nodosaurids, such as *Mymoorapelta* (Kirkland and Carpenter, 1994). In ventral view, the ischial and pubic peduncles are prominent and continuous with each other as in most ankylosaurians, such as *Jinyunpelta* (Zheng et al., 2018), but unlike the well-separated peduncles in *Gargoyleosaurus* and *Mymoorapelta*




Fig. 2. Photographs (left) and drawings (right) of the ankylosaurian ilium (IVPP V26052) from Yanji in lateral (A), dorsal (B), ventral (C), and medial (D) views. Abbreviations: ac, acetabulum; ip, ischial peduncle; pop, postacetabular process; pp, pubic peduncle; prp, preacetabular process; prr, preacetabular ridge; prs, preacetabular shelf.

(Carpenter et al., 2013). The anterior portion of the peduncles is extended more ventrally and is more prominent than the posterior portion. The peduncles bound the acetabulum medially, which is shallow, cup-like and imperforate, as in other ankylosaurians (Vickaryous et al., 2004). Generally, this ventral surface bears a well-marked rounded longitudinal preacetabular ridge in ankylosaurians (Fig. 2: prr), which starts from the anterolateral corner of the iliac blade, becomes gradually fainter posteriorly, and merges with the ilium body anterolateral to the acetabulum. This ridge forms the lateral edge of the preacetabular process in the anterior portion, making the iliac blade thicker laterally than medially anterior to the acetabulum (Coombs, 1979). The preacetabular ridge is straight in ventral view, as in *Zhejiangosaurus* (Lü et al., 2007b), *Gargoyleosaurus* (Carpenter et al., 2013), *Sauropelta* (Coombs, 1978), *Chuanqilong* (Han et al., 2014), *Anodontosaurus* (AMNH 5245) (Arbour and Currie, 2013), and *Jinyunpelta* (Zheng et al., 2018). In contrast, the ridge is well-curved in *Crichtonpelta* (Arbour and Currie, 2016; Lü et al., 2007a), and *Pinacosaurus* cf. *grangeri* from Shandong (Buffetaut, 1995). The preacetabular ridge separates a narrow preacetabular shelf (Fig. 2: prs) from the main body of the preacetabular process, as in other ankylosaurians (Carpenter et al., 2013; Lü et al., 2007a; Lü et al., 2007b). The preacetabular process becomes slightly narrower anteriorly but is still relatively wide, like those of *Taohelong* (Yang et al., 2013), *Talarurus* (Maleev, 1956), and *Jinyunpelta* (Zheng et al., 2018). The process tapers gradually in contrast to those of some other ankylosaurians, such as *Gargoyleosaurus* (Carpenter et al., 2013), and *Gastonia* (Kirkland, 1998), in which the process tapers abruptly anterior to the acetabulum. In medial view, the medial edge is almost horizontal.

5. Discussion

The incomplete preservation of IVPP V26052 makes body size estimation difficult. The mediolateral breadth of the preacetabular process is approximately 19.0 cm at the point where the preacetabular ridge merges into the lateral edge (Fig. 2C). Although the medial edge of IVPP V26052 is somewhat damaged, this is common in ankylosaurian ilia due to their thin medial edge as seen in *Dyoplosaurus* (Arbour et al., 2009; Parks, 1924), *Anodontosaurus* (AMNH 5245) (Arbour and Currie, 2013), and *Jinyunpelta* (ZMNH M8961) (Zheng, 2018). The preserved mediolateral breadth of the preacetabular process of *Jinyunpelta* (ZMNH M8961) is approximately 15.8 cm (Fig. 3) (Zheng, 2018). The whole anteroposterior length of the ilium of ZMNH M8961 is 90.0 cm, similar to *Polacanthus* (NHMUK PV R175) (Pereda-Suberbiola, 1994; Raven et al., 2020). The estimated body lengths are 4.5–5 m for *Jinyunpelta* (Zheng, 2018) and 5–7 m for *Polacanthus* (Pereda-Suberbiola, 1994). The medial edge of the ilium of *Jinyunpelta* (ZMNH M8961) is already very thin, suggesting that the original form of the medial edge is mostly preserved and thus the width of the preacetabular process would not exceed 19.0 cm. The breadth of the preacetabular process at the similar position of *Polacanthus* (is approximately 21.0 cm, which is broader than IVPP V26052. *Euoplocephalus* (AMNH 5409) and *Edmontonia longiceps* (CMN 8531) also share a broader preacetabular process (Arbour and Currie, 2013; Coombs, 1979), and both of them have longer ilia than *Jinyunpelta* or *Polacanthus* (Arbour and Currie, 2013; Carpenter, 1990). The estimated body length of *Euoplocephalus* and *Edmontonia longiceps* are 5–6 m and 6–7 m, respectively (Arbour and Mallon, 2017; Vickaryous et



Fig. 3. The left ilium from Yanji (IVPP V26052) (A) and the left ilium of *Jinyunpelta* (ZMNH M8961) (B) in ventral views.

al., 2004). The small-bodied *Struthiosaurus languedocensis* has a maximum width of the preacetabular process reaching 12.5 cm and an estimated body length ranging between 2.5 and 3 m (Garcia and Suberbiola, 2003). Based on these comparisons, we estimate the body length of IVPP V26052 to be around 5–5.5 m, longer than *Jinyunpelta* but shorter than *Polacanthus*, *Euoplocephalus*, or *Edmontonia*. The estimated body length of the largest ankylosaurians exceeds 7 m, as in *Ankylosaurus* (6–10 m) (Arbour and Mallon, 2017) and *Cedarpelta* (7.5–8.5 m) (Carpenter et al., 2001). Other large-bodied ankylosaurians with a body length around 5–6 m include *Zuul* and ZPAL MgD I/113 (Ankylosaurinae indet.) (Arbour and Mallon, 2017), *Borealopelta*, *Sauropelta* (Brown et al., 2017), and *Scolosaurus* (Penkalski and Blows, 2013). Small-to medium-sized ankylosaurians with a body length around 3–5 m include *Gargoyleosaurus* (3–3.5 m) (Kilbourne and Carpenter, 2005), *Hungarosaurus* (4–4.5 m) (Ösi et al., 2019; Ösi and Makádi, 2009), *Pinacosaurus grangeri* (5 m) (Maryańska, 1977), *Pinacosaurus mephistocephalus* (3 m) (Godefroit et al., 1999), PIN 614 (3.6 m) and MPC 100/1305 (3.9 m) (both cf. *Pinacosaurus*), *Dyoplosaurus* (3.95–4.53 m), and *Anodontosaurus* (>4 m) (Arbour and Mallon, 2017). The small-sized ankylosaurians are less than 3 m, such as *Struthiosaurus*, *Minmi*, *Mymoorapelta*, and *Kunbarrasaurus* (Kirkland and Carpenter, 1994; Garcia and Suberbiola, 2003; Leahey et al., 2015). Thus, given the lack of information indicating the precise ontogenetic stage of the specimen, the ankylosaurian specimen from the Longjing Formation (IVPP V26052) at least represents a moderate body-sized ankylosaurian dinosaur.

Ankylosaurian remains are very abundant in China and have been reported from the Jurassic of the Junggar Basin in Xinjiang Uyghur Autonomous Region (Augustin et al., 2020; Dong, 1993); the Lower Cretaceous of Yixian (Xu et al., 2001), the Lower to Upper Cretaceous of Beipiao (Dong, 2002; Lü et al., 2007a; Yang et al., 2017), the Lower Cretaceous of Fuxin (Amiot et al., 2010), the Lower Cretaceous of Lingyuan (Han et al., 2014), and the Lower Cretaceous of Kazuo (Ji et al., 2014) in Liaoning Province; the Lower Cretaceous of Yongjing in Gansu Province (Yang et al., 2013), the Lower Cretaceous of Urad Houqi (Arbour and Currie,

2016; Bohlin, 1953), the Lower and Upper Cretaceous of Alashan (Vickaryous et al., 2001; Young, 1935), and the Upper Cretaceous of Bayan Mandahu (Burns et al., 2011; Godefroit et al., 1999) in Inner Mongolia Autonomous Region; the Upper Cretaceous of Tianzhen (Barrett et al., 1998; Pang and Cheng, 1998), and Zuoyun (Jia, 2018) in Shanxi Province; the Upper Cretaceous of Laiyang in Shandong Province (Buffetaut, 1995); the Cretaceous of Ruyang in Henan Province (Xu et al., 2007); the Cretaceous of Lishui (Lü et al., 2007b), Dongyang (Chen et al., 2013) and Jinyun (Zheng et al., 2018) in Zhejiang Province (Fig. 4). Among them, the localities of Beipiao in Liaoning, Zuoyun in Shanxi, and Lishui, Dongyang, and Jinyun in Zhejiang have similar geological ages to the Longjing Formation (Chen et al., 2013; Yang et al., 2017; Jia, 2018; Zheng et al., 2018). Only two of them are from North China. Wang et al. (2020) reported a new ankylosaur *Sinankylosaurus zhuchengensis* from Upper Cretaceous of Zhucheng, Shandong Province, based on an incomplete right ilium (ZJZ-183). However, this holotype presents no ankylosaur diagnostic features. For instance, the putative acetabulum does not have cup-like structure, nor pubic or ischial peduncle. Moreover, there is no longitudinal preacetabular ridge in the ventral surface anterior to the putative acetabulum (Wang et al., 2020: Fig 2–3). For those reasons, we concluded that ZJZ-183 cannot be referred to an ankylosaur ilium and the diagnosis for ZJZ-183 as a new ankylosaur taxon is invalid. ‘*Sinankylosaurus zhuchengensis*’ is here considered as a nomen dubium, and not an ankylosaur.

The ilium described herein is the first evidence of ankylosaurian dinosaurs from Jilin Province, and is the easternmost known ankylosaurian occurrence in China at present (Fig. 4). Thus, this discovery extends the geographical range of ankylosaurians, and also enriches the middle Cretaceous ankylosaurian record, especially in North China. Unfortunately, the recovered portion prevents further referral of IVPP V26052 to any existing taxa, or to either the subclade of Nodosauridae or Ankylosauridae. The discovery demonstrates the potential of ongoing excavation in the Longjing Formation of Longshan Mountain to recover additional ankylosaurian remains (Zhang et al., 2018).



Fig. 4. Ankylosaurian fossil localities in China. Jurassic and Cretaceous localities are indicated by black squares and red triangles, respectively. (For interpretation of the references to colour in this figure legend, the reader is referred to the Web version of this article.)

6. Conclusions

The new ankylosaurian ilium (IVPP V26052), from the Albian–Cenomanian Longjing Formation of Yanji City, Jilin Province, northeastern China, represents the first definitive ankylosaurian dinosaur from Jilin Province, and the easternmost ankylosaurian occurrence in China. We estimated the body length of the individual to be around 5–5.5 m. Given the lack of precise ontogenetic stage information of the specimen, it at least represents a moderate body-sized ankylosaurian dinosaur.

Acknowledgements

We thank Lishi Xiang (IVPP), Sicai He (IVPP), and Yushan Shen (IVPP) for participating in the excavation of the specimen, Lishi Xiang for preparing the specimen. We thank the following people for access to and assistance at their respective institutions: Binghe Geng (IVPP), Mark Norell (AMNH), Carl Mehling (AMNH), Xiaochun Wu (CMN), Margaret Currie (CMN), Kieran Shepherd (CMN), Paul Barrett (NHMUK), and Susannah Maidment (NHMUK). This research was supported by the National Natural Science Foundation of China (Grant No. 41602019, 41688103), and the State Key Laboratory of Palaeobiology and Stratigraphy (Nanjing Institute of

Geology and Palaeontology, CAS) (Grant No. 163120). We thank the editor, Dr. Paul Barrett (NHMUK), and an anonymous reviewer for their helpful comments and suggestions on this manuscript.

References

- Amiot, R., Kusuhashi, N., Xu, X., Wang, Y., 2010. Isolated dinosaur teeth from the Lower Cretaceous Shahaai and Fuxin formations of northeastern China. *Journal of Asian Earth Sciences* 39, 347–358.
- Arbour, V.M., Currie, P.J., 2013. *Euoplocephalus tutus* and the diversity of ankylosaurid dinosaurs in the Late Cretaceous of Alberta, Canada, and Montana, USA. *PLoS One* 8, e62421.
- Arbour, V.M., Currie, P.J., 2016. Systematics, phylogeny and palaeobiogeography of the ankylosaurid dinosaurs. *Journal of Systematic Palaeontology* 14, 385–444.
- Arbour, V.M., Mallon, J.C., 2017. Unusual cranial and postcranial anatomy in the archetypal ankylosaur *Ankylosaurus magniventris*. *FACETS* 2, 764–794.
- Arbour, V.M., Burns, M.E., Sissons, R.L., 2009. A redescription of the ankylosaurid dinosaur *Dyoplosaurus acutosquameus* Parks, 1924 (Ornithischia: Ankylosauria) and a revision of the Genus. *Journal of Vertebrate Paleontology* 29, 1117–1135.
- Augustin, F.J., Matzke, A.T., Maisch, M.W., Pfretzschner, H.-U., 2020. First evidence of an ankylosaur (Dinosauria, Ornithischia) from the Jurassic Qigu Formation (Junggar Basin, NW China) and the early fossil record of Ankylosauria. *Geobios* 61, 1–10.
- Barrett, P.M., You, H.L., Upchurch, P., Burton, A.C., 1998. A new ankylosaurian dinosaur (Ornithischia: Ankylosauria) from the Upper Cretaceous of Shanxi Province, People's Republic of China. *Journal of Vertebrate Paleontology* 18, 376–384.
- Bohlin, B., 1953. Fossil reptiles from Mongolia and Kansu. *Sino-Swedish Expedition Publication* 37, 1–113.

- Brown, C.M., Henderson, D.M., Vinther, J., Fletcher, I., Sistiaga, A., Herrera, J., Summons, R.E., 2017. An exceptionally preserved three-dimensional armored dinosaur reveals insights into coloration and Cretaceous predator-prey dynamics. *Current Biology* 27, 2514–2521.e2513.
- Buffetaut, E., 1995. An ankylosaurid dinosaur from the Upper Cretaceous of Shandong (China). *Geological Magazine* 132, 683–692.
- Bureau of Geology and Mineral Resources of Jilin Province, 1988. *Regional Geology of Jilin Province*. Geological Publishing House, Beijing, pp. 224–258.
- Burns, M.E., Currie, P.J., Sissons, R.L., Arbour, V.M., 2011. Juvenile specimens of *Pinacosaurus grangeri* Gilmore, 1933 (Ornithischia: Ankylosauria) from the Late Cretaceous of China, with comments on the specific taxonomy of *Pinacosaurus*. *Cretaceous Research* 32, 174–186.
- Carpenter, K., 1990. Ankylosaur systematics: example using *Panoplosaurus* and *Edmontonia* (Ankylosauria: Nodosauridae). In: Carpenter, K., Currie, P.J. (Eds.), *Dinosaur systematics: perspectives and approaches*. Cambridge University Press, Cambridge, pp. 281–298.
- Carpenter, K., Kirkland, J.I., Burge, D., Bird, J., 2001. Disarticulated skull of a new primitive ankylosaurid from the Lower Cretaceous of Eastern Utah. In: Carpenter, K. (Ed.), *The armored dinosaurs*. Indiana University Press, Bloomington, pp. 211–238.
- Carpenter, K., DiCroce, T., Kinneer, B., Simon, R., 2013. Pelvis of *Gargoylesaurus* (Dinosauria: Ankylosauria) and the origin and evolution of the ankylosaur pelvis. *PLoS One* 8, e79887.
- Chen, R., Zheng, W., Azuma, Y., Shibata, M., Lou, T., Jin, Q., Jin, X., 2013. A new nodosaurid ankylosaur from the Chaochuan Formation of Dongyang, Zhejiang Province, China. *Acta Geologica Sinica (English Edition)* 87, 658–671.
- Coombs Jr., W.P., 1978. The families of the ornithischian dinosaur Order Ankylosauria. *Palaeontology* 21, 143–170.
- Coombs Jr., W.P., 1979. Osteology and myology of the hindlimb in the Ankylosauria (Reptilia, Ornithischia). *Journal of Paleontology* 53, 666–684.
- Cui, T., Ying, Z., Zhang, Y., 1994. Strata sequence of Late Mesozoic in Yanji Basin. *Petroleum Geology & Oilfield Development in Daqing* 13, 25–28.
- Dong, Z.-M., 1993. An ankylosaur (ornithischian dinosaur) from the Middle Jurassic of the Junggar Basin, China. *Vertebrata Palasiatica* 31, 257–266.
- Dong, Z.-M., 2002. A new armored dinosaur (Ankylosauria) from Beipiao Basin, Liaoning Province, Northeastern China. *Vertebrata Palasiatica* 40, 276–285.
- García, G., Suberbiola, X.P., 2003. A new species of *Struthiosaurus* (Dinosauria: Ankylosauria) from the Upper Cretaceous of Villeveyrac (southern France). *Journal of Vertebrate Paleontology* 23, 156–165.
- Godefroit, P., Pereda-Suberbiola, X., Li, H., Dong, Z.-M., 1999. A new species of the ankylosaurid dinosaur *Pinacosaurus* from the Late Cretaceous of Inner Mongolia (P. R. China). *Bulletin de l'Institut Royal des Sciences Naturelles de Belgique: Sciences de la Terre* 69, 17–36.
- Han, F., Zheng, W., Hu, D., Xu, X., Barrett, P.M., 2014. A new basal ankylosaurid (Dinosauria: Ornithischia) from the Lower Cretaceous Jiufotang Formation of Liaoning Province, China. *PLoS One* 9, e104551.
- Ji, S.-a., Zhang, L., Zhang, S., Zhang, L., Hang, S., 2014. Large-sized ankylosaur (Dinosauria) from the Lower Cretaceous Jiufotang Formation of Western Liaoning, China. *Acta Geologica Sinica - English Edition* 88, 1060–1065.
- Jia, L., 2018. A new ankylosaurid dinosaur from Zouyuan of Shanxi Province, China, Institute of Vertebrate Paleontology and Paleoanthropology, Chinese Academy of Sciences. University of Chinese Academy of Sciences, pp. 1–55.
- Jin, D., Zhang, J., Xu, X., Jin, C., Jin, F., Cai, Y., 2018. A preliminary report on the Yanji dinosaur fauna in Jilin. *Acta Palaeontologica Sinica* 57, 495–503.
- Kilbourne, B.M., Carpenter, K., 2005. Redescription of *Gargoylesaurus parkpinorum*, a polacanthid ankylosaur from the Upper Jurassic of Albany County, Wyoming. *Neues Jahrbuch für Geologie und Paläontologie - Abhandlungen* 237, 111–160.
- Kirkland, J.I., 1998. A polacanthine ankylosaur (Ornithischia: Dinosauria) from the Early Cretaceous (Barremian) of eastern Utah. *New Mexico Museum of Natural History and Science Bulletin* 14, 271–281.
- Kirkland, J.I., Carpenter, K., 1994. North America's first pre-Cretaceous ankylosaur (Dinosauria) from the Upper Jurassic Morrison Formation of western Colorado. *Brigham Young University Geology Studies* 40, 25–42.
- Leahey, L.G., Molnar, R.E., Carpenter, K., Witmer, L.M., Salisbury, S.W., 2015. Cranial osteology of the ankylosaurian dinosaur formerly known as *Minmi* sp. (Ornithischia: Thyreophora) from the Lower Cretaceous Allaru Mudstone of Richmond, Queensland, Australia. *PeerJ* 3, e1475.
- Li, X., Gong, G., 2008. Constraints from Fission Track Thermochronology for Geological age of Dalazi Formation in Jilin Province, China. *Atomic Energy Science and Technology* 42, 565–567.
- Lü, J., Ji, Q., Gao, Y., Li, Z., 2007a. A new species of the ankylosaurid dinosaur *Crichtonsaurus* (Ankylosauridae: Ankylosauria) from the Cretaceous of Liaoning Province, China. *Acta Geologica Sinica - English Edition* 81, 883–897.
- Lü, J., Jin, X., Sheng, Y., Li, Y., Wang, G., Azuma, Y., 2007b. New nodosaurid dinosaur from the Late Cretaceous of Lishui, Zhejiang Province, China. *Acta Geologica Sinica (English Edition)* 81, 344–350.
- Maleev, E.A., 1956. Armored dinosaurs of the Upper Cretaceous of Mongolia, Family Ankylosauridae. *Trudy Palaeontologicheskoi Instytuta. Akademiia Nauk SSSR* 62, 51–91.
- Maryńska, T., 1977. Ankylosauridae (Dinosauria) from Mongolia. *Palaeontologica Polonica* 37, 85–151.
- Matsukawa, M., Futakami, M., Lockley, M.G., Chen, P., Chen, J., Cao, Z., Bolotsky, U.L., 1995. Dinosaur footprints from the Lower Cretaceous of eastern Manchuria, northeastern China: Implications for the recognition of an ornithopod ichnofacies in East Asia. *PALAIOS* 10, 3–15.
- Nopcsa, F., 1915. Die dinosaurier der siebenbürgischen Landesteile Ungarns, vol. 23, pp. 1–26. *Mitt. Jb. K. Ungar. Geo. Reichsanst., Budapest*.
- Ósi, A., Botfalvai, G., Albert, G., Hajdu, Z., 2019. The dirty dozen: taxonomical and taphonomical overview of a unique ankylosaurian (Dinosauria: Ornithischia) assemblage from the Santonian Iharkút locality, Hungary. *Palaeobiodiversity and Palaeoenvironments* 99, 195–240.
- Ósi, A., Makádi, L., 2009. New remains of *Hungarosaurus tormai* (Ankylosauria, Dinosauria) from the Upper Cretaceous of Hungary: skeletal reconstruction and body mass estimation. *Paläontologische Zeitschrift* 83, 227–245.
- Osborn, H.F., 1923. Two Lower Cretaceous dinosaurs of Mongolia. *American Museum Novitates* 95, 1–10.
- Owen, R., 1842. Report on british fossil reptiles. part II. Report - British Association for the Advancement of Science 11, 60–204.
- Pang, Q.-Q., Cheng, Z.-W., 1998. A new ankylosaur of the Late Cretaceous Tianzhen, Shanxi. *Progress in Natural Science* 8, 326–334.
- Parks, W., 1924. *Dyoplosaurus acutosquameus*, a new genus and species of armoured dinosaur. *University of Toronto Studies* 18, 1–35.
- Penkalski, P., Blows, W.T., 2013. *Scolosaurus cutleri* (Ornithischia: Ankylosauria) from the Upper Cretaceous Dinosaur Park Formation of Alberta, Canada. *Canadian Journal of Earth Sciences* 50, 171–182.
- Pereda-Suberbiola, J., 1994. *Polacanthus* (Ornithischia, Ankylosauria), a transatlantic armoured dinosaur from the Early Cretaceous of Europe and North America. *Palaeontographica Abteilung A Palaeozoologie-Stratigraphie* 232, 133–159.
- Raven, T.J., Barrett, P.M., Pond, S.B., Maidment, S.C.R., 2020. Osteology and taxonomy of British Wealden Supergroup (Berriasian–aptian) ankylosaurs (Ornithischia, Ankylosauria). *Journal of Vertebrate Paleontology* 40, e1826956.
- Seeley, H.G., 1887. On the classification of the fossil animals commonly named Dinosauria. *Proceedings of the Royal Society of London* 43, 165–171.
- Sun, G., Zheng, S., 2000. New proposal on division and correlation of Mesozoic from northeastern China. *Journal of Stratigraphy* 24, 60–64.
- Vickaryous, M.K., Maryńska, T., Weishampel, D.B., 2004. Ankylosauria. In: Weishampel, D.B., Dodson, P., Osmólska, H. (Eds.), *The Dinosauria*, second ed. University of California Press, Berkeley, pp. 363–392.
- Vickaryous, M.K., Russell, A.P., Currie, P.J., Zhao, X.-J., 2001. A new ankylosaurid (Dinosauria: Ankylosauria) from the Lower Cretaceous of China, with comments on ankylosaurian relationships. *Canadian Journal of Earth Sciences* 38, 1767–1780.
- Wang, K., Zhang, Y., Chen, J., Chen, S., Wang, P., 2020. A new ankylosaurian from the Late Cretaceous strata of Zhucheng, Shandong Province. *Geological Bulletin of China* 39, 958–962.
- Xing, L., Lockley, M.G., Kim, K.S., Klein, H., Matsukawa, M., Lim, J.D., Persons, W.S., Xu, X., 2017. Mid-Cretaceous dinosaur track assemblage from the Tongfosi Formation of China: Comparison with the track assemblage of South Korea. *Cretaceous Research* 74, 155–164.
- Xu, X., Wang, X.-L., You, H.-L., 2001. A juvenile ankylosaur from China. *Naturwissenschaften* 88, 297–300.
- Xu, L., Lü, J.-C., Zhang, X.-L., Jia, S.-H., Hu, W.-Y., Zhang, J.-M., Wu, Y.-H., Ji, Q., 2007. New nodosaurid ankylosaur from the Cretaceous of Ruyang, Henan Province. *Acta Geologica Sinica* 81, 433–438.
- Yang, J.-T., You, H.-L., Li, D.-Q., Kong, D.-L., 2013. First discovery of polacanthine ankylosaur dinosaur in Asia. *Vertebrata Palasiatica* 51, 265–277.
- Yang, J., You, H., Xie, L., Zhou, H., 2017. A new specimen of *Crichtonpelta benxiensis* (Dinosauria: Ankylosaurinae) from the Mid-Cretaceous of Liaoning Province, China. *Acta Geologica Sinica (English Edition)* 91, 781–790.
- Young, C.C., 1935. On a new nodosaurid from Ninghsia. *Geological Survey of China, Beijing*, pp. 1–28.
- Zhang, L., Jin, C., Yin, Q., Huyskens, M., Jin, D., Zhang, J., Jin, F., Xu, X., 2018. A new dinosaur fossil locality of mid-Cretaceous age in northeastern China. In: Farke, A., MacKenzie, A., Miller-Camp, J. (Eds.), *The 78th Annual Meeting of the Society of Vertebrate Paleontology, Albuquerque, New Mexico, U.S.A.*, p. 247.
- Zheng, W., 2018. New ankylosaurid material from the Cretaceous of China and the early evolution of Ankylosauridae, Institute of Vertebrate Paleontology and Paleoanthropology, Chinese Academy of Sciences. University of Chinese Academy of Sciences, pp. 1–290.
- Zheng, W., Jin, X., Azuma, Y., Wang, Q., Miyata, K., Xu, X., 2018. The most basal ankylosaurine dinosaur from the Albian–Cenomanian of China, with implications for the evolution of the tail club. *Scientific Reports* 8, 3711.
- Zhong, Y., Wang, Y., Jia, B., Wang, M., Hu, L., Pan, Y., 2020. A potential terrestrial Albian–Cenomanian boundary in the Yanji Basin, Northeast China. *Palaeogeography, Palaeoclimatology, Palaeoecology* 110088.