



Historical Biology

An International Journal of Paleobiology

ISSN: 0891-2963 (Print) 1029-2381 (Online) Journal homepage: <http://www.tandfonline.com/loi/ghbi20>

The second mamenchisaurid dinosaur from the Middle Jurassic of Eastern China

Xin-Xin Ren, Jian-Dong Huang & Hai-Lu You

To cite this article: Xin-Xin Ren, Jian-Dong Huang & Hai-Lu You (2018): The second mamenchisaurid dinosaur from the Middle Jurassic of Eastern China, *Historical Biology*, DOI: [10.1080/08912963.2018.1515935](https://doi.org/10.1080/08912963.2018.1515935)

To link to this article: <https://doi.org/10.1080/08912963.2018.1515935>



Published online: 19 Sep 2018.



Submit your article to this journal [↗](#)



Article views: 53



View Crossmark data [↗](#)

ARTICLE



The second mamenchisaurid dinosaur from the Middle Jurassic of Eastern China

Xin-Xin Ren^{a,b,c,d}, Jian-Dong Huang^d and Hai-Lu You^{a,b,c}

^aKey Laboratory of Vertebrate Evolution and Human Origins of Chinese Academy of Sciences, Institute of Vertebrate Paleontology and Paleoanthropology, Chinese Academy of Sciences, Beijing, China; ^bCAS Center for Excellence in Life and Palaeoenvironment, Beijing, China; ^cUniversity of the Chinese Academy of Sciences, Beijing China; ^dAnhui Geological Museum, Hefei, China

ABSTRACT

A new mamenchisaurid dinosaur, *Anhuilong diboensis* gen. et sp. nov. from the Middle Jurassic of Eastern China is reported here. The holotype consists of complete left humerus, ulna and radius of an individual. Comparative study and cladistic analysis shows this new taxon belongs to Mamenchisauridae and bears a unique combination of characters, such as low ratios of the average of the greatest widths of the proximal end, mid-shaft and distal end of the humerus/length of the humerus, total length of ulna to humerus and total length of radius to humerus; the lateral edge of the deltopectoral crest directs caudolaterally, the lateral accessory condyle on the craniodistal edge of humerus is more robust than the medial one, and the cross-sectional shape of the ulna at mid-shaft is elliptical with highest ratio of transverse to craniocaudal diameter among mamenchisaurids. Phylogenetically, *Anhuilong* is the sister taxon of *Huangshanlong*, and with *Omeisaurus* they together form the sister clade to all other members of Mamenchisauridae. Including *Huangshanlong*, two mamenchisaurids have been found in eastern China, and indicates that Mamenchisauridae was already a diverse sauropod clade in China by the Middle Jurassic.

ARTICLE HISTORY

Received 12 June 2018
Accepted 22 August 2018

KEYWORDS

Huangshan; Eastern China; Middle Jurassic; Hongqin Formation; Mamenchisauridae; *Anhuilong*

Introduction

Mamenchisauridae (Young and Chao 1972) is the dominant sauropod clade in the Middle and Late Jurassic of East Asia (Lü et al. 2008; Upchurch et al. 2004; Mannion et al. 2011; Suteethorn 2012; Xing et al. 2015b). It contains multiple genera, although many of them are based on fragmentary and undiagnosable materials (Young 1939, 1954, 1958; Young and Chao 1972; Hou et al. 1976; He et al. 1988, 1996; Zhao et al. 1993; Russell and Zheng 1993; Zhang et al. 1998; Tang et al. 2001; Li 1997; Fang et al. 2000, 2004; Ouyang et al. 2002; Lü et al. 2008; Jiang et al. 2011; Wu et al. 2013; Huang et al. 2014; Xing et al. 2015b). This clade has been referred to Diplodocidae, Bothrosauropodoidea, and 'Euhelopodidae', the current consensus regards the group as an early-diverging clade within Eusauropoda. Still, the relationships among the members of the clade are unstable, even the validity of Mamenchisauridae is debatable (Li and Cai 1997; Upchurch et al. 2004; McIntosh 1990; Wilson and Sereno 1998; Xing et al. 2015b). Additionally, Mamenchisauridae may not be endemic in East Asia (Suteethorn 2012; Mannion 2013). Sauropod fossil records are sparse in the Middle Jurassic (Upchurch et al. 2004; Barrett and Upchurch 2005; Remes et al. 2009), and mamenchisaurids from the Middle Jurassic of China provide valuable opportunity to understand sauropod diversity and phylogeny in this period.

Here we report a new mamenchisaurid sauropod, *Anhuilong diboensis* gen. et sp. nov., based on a left humerus, ulna and radius from the Middle Jurassic Hongqin Formation of Anhui Province, Eastern China (Figure 1). This formation has previously yielded the mamenchisaurid *Huangshanlong* (Huang et al. 2014). Comparative study and cladistic analysis shows the new taxon is different from *Huangshanlong*.

Anhuilong, *Huangshanlong*, and *Omeisaurus* form the sister clade of all other known mamenchisaurids.

Geological setting

Many continental basins developed in Eastern China during the Mesozoic. Tunxi Basin is one of them located in the midland, pertaining to a small intermountain basin during the Mesozoic. In this basin, the red beds are well exposed, and the outcrops are generally continuous. It is a representative basin on stratigraphic research in the eastern part of China (Yu et al. 2001). The geotectonic position of Tunxi Basin belongs to the eastern segment of Jingnan intracontinental orogenic belt, southeast of Yangzi Block (Jiang et al. 2016). Since the Indosinian Movement, this area is exposed above the sea level as a result of the crustal rising. Different types of continental basins were chronologically overlapped during different tectonic phases. Tunxi Basin is dominated by fluvial deposits with volcanic deposits in Jurassic (Ren et al. 2015 and 2017).

The new genus comes from purple sandstones of the Hongqin Formation, interbedded with purple shales. The Hongqin Formation was originally regarded to be of Middle Jurassic age according to the study of invertebrate (Bureau of Geology and Mineral Resources of Anhui Province 332 Geological team 1971, unpublished).

Systematic paleontology

Dinosauria Owen (1842)
Saurischia Seeley (1887)




Figure 1. Geographic and geologic map showing the location of *Anhuilong diboensis* gen. et sp. nov. (indicated by the red star and dinosaur silhouette) and generalized stratigraphic section of Jurassic of Tunxi Basin, modified from Ren et al. (2017).

Sauropodomorpha Huene (1932)
 Sauropoda Marsh (1878)
 Eusauropoda Upchurch (1995)
 Mamenchisauridae Young and Chao (1972)
Anhuilong diboensis gen. et sp. nov. (Figures. 2–4)

Etymology

The generic name refers to Anhui Province, where the holotype was found; ‘long’ means dragon in Chinese Pinyin. The specific name refers to the locality where the holotype was repositied.

Holotype

Anhui Geological Museum: AGB 5822. Associated and complete left humerus, ulna and radius. Reposited in Anhui Geological Museum, Hefei, Anhui Province, China.

Diagnosis

A mamenchisaurid possessing the following unique combination of character states (autapomorphies are marked by *):




Figure 2. Left humerus of *Anhuilong diboensis* gen. et sp. nov. A, cranial; B, medial; C, caudal; D, lateral; E, proximal; F, cross section in proximal (the cross section near the narrowest area); G, distal views (the cranial is upward). Scale bar equals 10cm. Abbreviations: cod, caudolateral bulge of the deltopectoral crest; dpc, deltopectoral crest; map, medial accessory process; mep, medial part of proximal surface; mip, middle part of proximal surface; lap, lateral accessory process; lp, lateral part of proximal surface; plc, proximolateral corner; plp, proximolateral process.




Figure 3. Left ulna of *Anhuilong diboensis* gen. et sp. nov. A, cranial; B, medial; C, caudal; D, lateral; E, proximal (the cranial is upward); F, distal views (the cranial is upward); G, reconstruction of left ulna and radius (gray represents ulna, blue represents radius; i, craniolateral; ii, proximal; iii, distal views). Scale bar equals 10cm. Abbreviations: clp, craniolateral process; cmcu, craniomedial convex of lower part of ulna; cmp, craniomedial process.




Figure 4. Left radius of *Anhuilong diboensis* gen. et sp. nov. A, cranial; B, medial; C, caudal; D, lateral; E, proximal (the cranial is upward); F, the distal views (the cranial is upward); G, reconstruction of left ulna and radius (gray represents ulna, blue represents radius; i, craniolateral; ii, proximal; iii, distal views). Scale bar equals 10cm. Abbreviations: cmcr, caudomedial convex of lower part of radius; lr, longitudinal ridge.

lateral edge of deltopectoral crest turning caudolaterally; lateral accessory process of humerus more robust than medial one; ratio of total length of radius to humerus is 0.50 (the lowest value among mamenchisaurids; *Huangshanlong*: 0.58)*; ratio of total length of ulna to humerus is 0.56 (the lowest value among mamenchisaurids; *Huangshanlong*: 0.67)*; elliptical cross-sectional shape of ulna at mid-shaft (*Huangshanlong* is circular), nearly perpendicular orientation of proximal end of ulna relative to long axis of ulnar shaft.

Locality and horizon

The specimen was excavated at Jimushan of Hengguan Village, Wangcun Town, Shexian County, Huangshan City, Anhui Province, Eastern China. The quarry is from the lower part of the Middle Jurassic Hongqin Formation. The bones were isolatedly found in a small area with a massive to finely laminated red siltstone containing some carbonate in its

matrix. The siltstone layer is several meters thick and yielded the sauropod remains in its upper half. The sedimentological and petrologic researches indicate that the fossils were deposited in the floodplain environment (Wang et al. 2006).

Description

The left humerus is well preserved (Figure 2; see Table 1 for measurement). The outline is similar to other mamenchisaurids, but it looks a little more slender than other taxa of Mamenchisauridae with the lowest RI value (RI, the average of the greatest widths of the proximal end, mid-shaft and distal end of humerus/length of humerus). In cranial view, the proximal and distal portions expand gradually towards both ends, giving the proximal end a fan-like shape in cranial view. The proximal width is 36% of the total length of the humerus, resembling that in *Shunosaurus lii*, *Barapasaurus*, *Patagosaurus*, *Camarasaurus*, *Alamosaurus* and *Turiasaurus*

Table 1. Measurements of the left humerus (in cm).

Total length	105
Transverse width of proximal end	38
Craniocaudal width of proximal end (across articular head)	17
Distance from the proximal end to the lateral margin of the deltopectoral crest	30
Transverse width of mid-shaft	14
Craniocaudal width of mid-shaft	10
Transverse width of distal end	30
Maximum craniocaudal width of the distal end (across the two condyles)	20
RI	0.26

RI: the average of the greatest widths of the proximal end, mid-shaft and distal end of humerus/length of humerus (after Wilson and Upchurch 2003)

(Dong et al. 1983; Bonaparte 1986; McIntosh et al. 1996; Lehman and Coulson 2002; Royo-Torres et al. 2006; Bandyopadhyay et al. 2010). By contrast, the ratio in *Huangshanlong* and other mamenchisaurids is approximately 41% (He et al. 1998; Fang et al. 2000; Tang et al. 2001; Ouyang and Ye 2002; Huang et al. 2014). A cranial projection overhangs on the middle of the proximal end, which is 2 cm in height, 28 cm in width. This cranial projection is weak in early-diverging sauropods and *Omeisaurus* spp. (Cooper 1981; Dong et al. 1983; He et al. 1988; Tang et al. 2001). The humeral head is located at the middle of the proximal end, and extends caudally in lateral view. The shape of the humeral head is close to an isosceles triangle, the base of which is 1.9 cm and the height is 0.9 cm. In proximal view, the proximal surface of humerus is divided into three planes, the lateral, middle, and medial (Figure 2, lp; mip; mep). The presence of these ‘three planes’ is common in mamenchisaurids, *Turiasaurus*, and some derived neosauropods forms such as *Tornieria* and *Saltasaurus* (He et al. 1988; Fang et al. 2000; Tang et al. 2001; Ouyang and Ye 2002; Remes 2006; Royo-

Torres et al. 2006). Supracoracoideus tuberculum is lacking on the proximolateral portion of the humerus as in some neosauropods such as *Suuwassea*, *Opisthocoelicaudia*, *Isisaurus* and *Saltasaurus* (Royo-Torres et al. 2006; Harris 2006a; Whitlock and Harris 2010).

The length of the deltopectoral crest/total length of humerus is about 0.44 (*Huangshanlong* is 0.43; *Omeisaurus tianfuensis* is 0.44). It is oriented craniolaterally, which differs from the conditions in other early diverging eusauropods (e.g. *Shunosaurus lii*, *Omeisaurus tianfuensis*, *Mamenchisaurus youngi*, *Chuanjiesaurus*, *Barapasaurus* and *Patagosaurus*). The distal half of the deltopectoral crest is 9 cm in length and the width is 2 cm. The caudolateral bulging of the deltopectoral crest is elliptical, the length of the major axis 10 cm and the minor axis 4 cm (Figure 2, cod). The cross section of the mid-shaft is ovoid, similar to those in *Vulcanodon*, *Shunosaurus lii*, *Omeisaurus tianfuensis* and *Huangshanlong*. The ratio of the width of the cross section to the total length of the humerus is 0.13, and this makes the morphology of *Anhuilong* particularly gracile considering the total length of the shaft (*Huangshanlong* is 0.16; *Omeisaurus tianfuensis* is 0.17). Despite the ovoid morphology of the cross section. The distal articular surface is deflected to the lateral and the angle between the axis of the distal articular surface and the proximal one is 25 degrees.

The distal end is quadrilateral, and the two condyles are convex slightly with coarse surfaces. There are two small accessory processes on the craniodistal edge of the humerus, as in other mamenchisaurids, *Patagosaurus*, *Apatosaurus*, and *Suuwassea*. Moreover, the lateral accessory process of *Anhuilong* is more robust than the medial one, which is different from other members of Mamenchisauridae (Figure 5). Both of the processes are nearly triangular. There is a shallow groove on the caudodistal portion of the humerus. The two condyles are semi-round in caudal view.




Figure 5. Comparison of forelimb of *Anhuilong*, *Huangshanlong* and *Omeisaurus tianfuensis*. A, left humerus of *Anhuilong* (from top to bottom: proximal; cranial; ventral views); B, right (left in reverse) humerus of *Huangshanlong* (from top to bottom: proximal; cranial; ventral views); C, left humerus of *Omeisaurus tianfuensis* (from top to bottom: proximal; cranial; ventral views); D, left ulna of *Anhuilong* (from top to bottom: proximal; lateral; ventral views); E, right (left in reverse) ulna of *Huangshanlong* (from top to bottom: proximal; lateral; ventral views); F, left ulna of *Omeisaurus tianfuensis* (from top to bottom: proximal; cranial; ventral views); G, left radius of *Anhuilong* (from top to bottom: proximal view; medial view; ventral view); H, right (left in reverse) radius of *Huangshanlong* (from top to bottom: proximal view; medial view; ventral view); I, left radius of *Omeisaurus tianfuensis* (from top to bottom: proximal view; medial view; ventral view); Abbreviation: cap, caudal process; clp, craniolateral process; cmcr, caudomedial convex of lower part of radius; cmcu, craniomedial convex of lower part of ulna; cmp, craniomedial process; dpc, deltopectoral crest; lap, lateral accessory process; map, medial accessory process; mip, middle part of proximal surface; lap, lateral accessory process; plc, proximolateral corner; plp, proximolateral process.

The left ulna is well preserved (Figure 3; see Table 2 for measurement). The ulna is longer than the radius. The proximal end is triradiate, similar to that in other sauropods. The cranialateral and cranio-medial processes are prominent and robust, which together in proximal view make up a 'L' shape and form a deep cranial groove that receives the proximal end of the radius. The angle between the two processes is 85 degrees, close to those in *Omeisaurus tianfuensis* (about 85 degrees) and *Huangshanlong* (about 75 degrees). The cranio-medial process is more developed than the cranialateral process. The length of the cranio-medial process is 1.5 times that of the cranialateral process (the distal end of the cranio-medial process is not well preserved), that in *Huangshanlong* is 1.7, and *Omeisaurus tianfuensis* is about 1.0. The caudal process is weakly developed, and the olecranon process is also weak. Along the cranialateral process, the maximum length is 31% of the total length of the ulna, and along the cranio-medial process, the maximum length is 45% (*Huangshanlong* is 40% and 44% respectively; *Omeisaurus tianfuensis* is 21% and 21% respectively). The middle part of the cranio-medial process is penetrated by a cranio-caudal groove, differing from the condition in other mamenchisaurids (e.g. *Omeisaurus tianfuensis*, *O. maodianus*, *Mamenchisaurus youngi*, *Chuanjiesaurus*, and *Huangshanlong*).

The 'L' shape of the proximal surface is transformed into an elliptical cross section at mid-shaft, resembling to the condition in *Huangshanlong*, *Vulcanodon*, *Chuanjiesaurus*, and *Rapetosaurus*, but the majority of sauropods have a circular cross section, such as in *Mamenchisaurus youngi*. On the cranialateral surface of the proximal portion, there is a raised coarse area that matches a similar area on the surface of the radius (Figure 3, rca).

The distal surface of the ulna is oval, and the cranial part is a little flat (the cranial is not preserved very well); in contrast, the shape of the corresponding region in *Omeisaurus tianfuensis* is more compressed, and those of *Mamenchisaurus youngi* and *Chuanjiesaurus* share a subquadrilateral shape. The maximum length of the distal end is 21% of the total length, similar to that in *Huangshanlong* (0.24). The distal portion of the cranio-medial surface is mildly convex where it received the caudomedial surface of the distal end of the radius. *Huangshanlong*, *Vulcanodon*, *Omeisaurus tianfuensis*, *Mamenchisaurus youngi*, and *Chuanjiesaurus* also share this morphology (Figure 3, cmcu). The surface of the distal end is nearly flat, with the center a little convex. The maximum angle between the distal surface and long axis of the shaft is 78 degrees, nearly similar to *Huangshanlong*, *Omeisaurus tianfuensis*, *Apatosaurus*, and *Amargasaurus* (Huang et al. 2014; He et al. 1998; Upchurch et al. 2004; Salgado and Bonaparte 1991). And still, it is larger than the early-diverging sauropods (e.g. *Vulcanodon* and *Shunosaurus lii*), *Barapasaurus*, *Alamosaurus*, *Opisthocoelicaudia*, *Lirainosaurus* (Borsuk-Bialynicka 1977; Cooper 1981; Dong et al. 1983; Company et al. 2009). Moreover, as an autapomorphy, the total length of the ulna is 56% of the total length of the humerus, which is the lowest value among mamenchisaurids (Table 3).

Table 2. Measurements of the left ulna (in cm).

Total length	59
Width of the proximal end (from the caudal surface to the tip of the cranio-medial process)	26
Width of the proximal end (from the caudal surface to the tip of the cranialateral process)	18
Transverse width of mid-shaft	11
Cranio-caudal width of mid-shaft	5.5
Transverse width of distal end	14
Maximum cranio-caudal width of the distal end	11

Table 3. Ratios of total length of ulna/total length of humerus (U/H) and total length of radius/total length of humerus (R/H) in mamenchisaurids. '-': not applicable.

Taxa	U/H	R/H
<i>Omeisaurus jiaoi</i>	0.75	0.71
<i>Omeisaurus maodianus</i>	0.62	0.65
<i>Omeisaurus tianfuensis</i>	-	0.70
<i>Mamenchisaurus anyuensis</i>	0.76	0.69
<i>Mamenchisaurus constructus</i>	-	-
<i>Mamenchisaurus fuxiensis</i>	-	-
<i>Mamenchisaurus hochuanensis</i>	-	-
<i>Mamenchisaurus youngi</i>	0.69	0.65
<i>Mamenchisaurus sinocanadorum</i>	-	-
<i>Mamenchisaurus jingyanensis</i>	0.66	0.60
<i>Eomamenchisaurus yuanmouensis</i>	-	-
<i>Chuanjiesaurus</i>	0.68	0.62
<i>Qijianglong</i>	-	-
<i>Huangshanlong</i>	0.67	0.58
<i>Xinjiangtitan</i>	-	-

The left radius is well-preserved and complete (Figure 4; see Table 4 for measurement). Proximally, the maximum width is 26% of the total length. The proximal end is nearly quadrilateral with a broad medial surface that met the cranio-medial process of ulna. The quadrilateral shape of the proximal end is similar to that in *Apatosaurus* (Upchurch et al. 2004), and different from the oval shape of that in early-diverging sauropods (e.g. *Vulcanodon*, *Shunosaurus lii*), some mamenchisaurids, *Barapasaurus*, and *Camarasaurus* (Dong et al. 1983; Cooper 1984; He et al. 1996; McIntosh et al. 1996; Tang et al. 2001; Bandyopadhyay et al. 2010). The surface of the proximal end is nearly flat with a slightly concave area in the center, and there is a bump cranio-medially, a condition also present in *Huangshanlong*, *Omeisaurus tianfuensis*, *Patagosaurus*, *Camarasaurus* and *Saltasaurus* (Bonaparte 1986; He et al. 1988; Powell 1992; McIntosh et al. 1996; Huang et al. 2014). The mid-shaft of the radius is elliptical in cross section. A prominent longitudinal ridge occurs on the proximal half of the radius and it is slightly twisted (Figure 4, lr). In lateral view, the caudal outline is

Table 4. Measurements of the left radius (in cm).

Total length	53
Width of the proximal end	10.5
Cranio-caudal width of the proximal end	14
Transverse width of mid-shaft	6
Cranio-caudal width of mid-shaft	8.5
Transverse width of distal end	10
Maximum cranio-caudal width of the distal end	15.5

prominently incurve. The surface of the proximal end is nearly perpendicular to the axis of the shaft. This condition is similar to *Omeisaurus tianfuensis*, *Patagosaurus*, and *Camarasaurus* (Bonaparte 1986; He et al. 1988; McIntosh et al. 1996).

The distal surface of the radius is flat, with an elliptical shape. A prominent elongated convex area exists on the caudomedial surface of distal portion. The distal radial condyle with respect to the long axis of shaft is nearly perpendicular (the maximum angle is less than 20 degrees), which is similar to most of sauropods (e.g. *Vulcanodon*, *Shunosaurus lii*, mamenchisaurids, *Apatosaurus*, and *Alamosaurus*).

Phylogenetic analysis

A phylogenetic analysis was conducted to assess the affinities of *Anhuilong diboensis* within Sauropoda (Figure 6). A maximum parsimony analysis used the data set of Xing et al. (2015b), with 344 original characters plus eight new forelimb characters (See the Supplementary Data). New characters based on descriptions and revisions (e.g.

Upchurch 1998; Upchurch et al. 2004; Wilson and Sereno 1998; Huang et al. 2014; Mannion et al. 2013; Wilson and Upchurch 2003), and entirely novel characters presented on the appendices, based on our personal observations and an extensive review of the literature. We did not exclude characters based on a priori assumptions about their level of homoplasy. Where possible, we tried to quantify, or at least more precisely define characters and state boundaries to remove ambiguity (see some similar attempts by Harris 2006). Both *Anhuilong* and *Huangshanlong* were added to the matrix. All characters were treated as unordered. The complete character list, including references, as well as our MESQUITE versions of the data matrices. The matrix was subjected to a heuristic search in TNT v. 1.5 (Goloboff et al. 2016), with multiple TBR + TBR search strategy (1000 replicated of trees, random addition sequence, tree bisection reconnection branch swapping algorithm, ten trees saved per replicate).

A strict consensus of all 450 most parsimonious trees (tree length = 1103; consistency index = 0.397; retention index = 0.638;) supports a monophyletic Mamenchisauridae. *Anhuilong* is recovered as the sister taxon of *Huangshanlong*,



Figure 6. Strict consensus of 450 MPTs (TL = 1104) from phylogenetic analysis (46 taxa, 352 characters). The data matrix follows Xing et al. (2015), with the addition of 8 character codings for *Anhuilong diboensis* gen. et sp. nov. (see Appendix 1).

and they together are closely related to *Omeisaurus tianfuensis* than to the clade including all other mamenchisaurids. Xing et al. (2015b) suggested that *O. maoianus* and *O. tianfuensis* are likely not congeneric, and *O. maoianus* is more closely related to *Mamenchisaurus*. Our phylogenetic analysis also supports this result. For this reason, all the statements of *Omeisaurus* just represents *O. tianfuensis* in this paper.

The mamenchisaurid clade is supported by ten unambiguous synapomorphy ('0' to '1' for character 64; 100; 102; 144; 146; 148; 307; 337; 347; '4' to '6' for character 105). *Anhuilong*, *Huangshanlong* and other mamenchisaurids clade is supported by the appearance of two cranial facing accessory processes on the craniodistal edge of humerus (character 347). The (*Anhuilong*+ *Huangshanlong* + *Omeisaurus*) clade is supported by two synapomorphies ('0' to '1' for character 238, 241): the stout ulna with high proximal breadth/proximodistal length ratio (character 238); radius distal/mid-shaft breadth ratio of 1.5–1.9 (character 241); Moreover, the sister-group relationship between *Anhuilong* and *Huangshanlong* is supported by one unambiguous character change ('1' to '0' for characters 349): the average of the greatest widths of the proximal end, mid-shaft and distal end/total length of humerus less than 0.27 (character 349).

Discussion

Though excavated in the same area, an analysis of morphology suggests that *Huangshanlong* (Huang et al. 2014) is different from *Anhuilong*. First of all, the shape of the humerus is stout compared to the comparatively gracile humerus of *Anhuilong*. Proximally, lateral and proximal humeral surfaces of *Huangshanlong* meet each other at an abrupt angle to produce a squared proximal end, making the proximolateral end a prominent process. On the contrary, the surfaces of *Anhuilong* merge smoothly with each other to produce a transversely rounded proximal end. The deltopectoral crest of *Huangshanlong* is much more robust than *Anhuilong*. The lateral humeral edge of *Huangshanlong* extends cranio-laterally, whereas, the lateral edge of the deltopectoral crest turns caudolateral in *Anhuilong*. Both *Huangshanlong* and *Anhuilong* preserve lateral and medial accessory process, as most mamenchisaurids do. The radial accessory process of *Huangshanlong* is more robust than the ulnar counterpart like other mamenchisaurids, while *Anhuilong* possesses the opposite condition. Many ridges exist on the ulna of *Huangshanlong*, while the ulna of *Anhuilong* has smooth texture. Moreover, the ratio of the ulnar length to the humeral length of *Huangshanlong* is much larger than *Anhuilong* (*Huangshanlong* is 0.67; *Anhuilong* is 0.56), also indicating an interesting fact that the antebrachium of *Anhuilong* is relatively short among mamenchisaurids.

Anhuilong also possesses some autapomorphies such as lowest ratio of ulnar length to the total length of humerus and the lowest ratio of radial length to the total length of humerus among mamenchisaurids. The ratio of ulnar length to total length of humerus is 0.56, which is the lowest value among all the taxa of Mamenchisauridae at present. This ratio

values on other taxa among the matrix are all about 0.60–0.80. And still, the ratio of radial length to total length of humerus is 0.50, also the lowest value among all mamenchisaurids so far, the values of other taxa are about 0.55–0.75. The functional implications of such a short antebrachium merits exploration by future studies.

Conclusions

The new genus *Anhuilong* shares characters with Mamenchisauridae. Our study confirms previous assignment of *Huangshanlong* as a member of Mamenchisauridae. Within this clade, the new genus is the sister taxon of *Huangshanlong*, and they together are more closely related to *Omeisaurus* than to the clade including all other mamenchisaurids.

Anhuilong enriches the diversity of early branching sauropods and provides additional information to help understand the evolutionary history of sauropods in Eastern China. This discovery indicates that Mamenchisauridae was already a diverse endemic sauropod lineage during the Middle Jurassic of Asia, and the interrelationships within mamenchisaurids need to be further explored.

Acknowledgments

The materials are collected by Han Li-Gang (deceased), from the Anhui Institute of Archaeology, to whom we express our sincerely respect. At the same time, we also appreciate the original collection unit of the holotype, the Cultural Relics Bureau of Shexian County (Shexian Museum).

Disclosure statement

No potential conflict of interest was reported by the authors.

Funding

This work was supported by the National Natural Science Foundation of China [41472020,41688103]; Non-Profit Sector Project, Ministry of Land and Resources of China [201511054].

ORCID

Hai-Lu You  <http://orcid.org/0000-0003-2203-6461>

References

- Bandyopadhyay S, Gillette DD, Ray S, Sengupta DP. 2010. Osteology of *barapasaurus tagorei* (dinosauria: sauropoda) from the jurassic of india. *Palaeontology*. 53:533–569.
- Barret PM, Upchurch P. 2005. Sauropodomorph diversity through time: possible macroevolutionary and palaeoecological implications. In: Curry-Rogers KA, Wilson JA, editors. *Sauropod evolution and paleobiology*. Berkeley: University of California Press; p. 125–156.
- Bonaparte JF. 1986. Les Dinosauriens (Carnosauriens, Allosauriens, Sauropodes, Cétiosauriens) du Jurassique Moyen de Cerro Cóndor (Chubut, Argentine). *Annales de Paléontologie*. 72:325–386.
- Borsuk-Bialynicka M. 1977. A new camarasaurid sauropod *Opisthocoelicaudia skarzynskii* gen. n., sp. n. from the Upper Cretaceous of Mongolia. *Palaeont. Pol.* 37:5–64.

- Bureau of Geology and Mineral Resources of Anhui Province 332 Geological team. 1971. unpublished. Qimen map H-50-22, Tunxi map H-50-23, 1:200 000 Regional Geological Mineral Survey Report.
- Casanovas ML, Santafé JV, Sanz JL. 2001. *Losillasaurus giganteus*, a new sauropod from the transitional jurassic-cretaceous of the los serranos basin (valencia, spain). *Paleontologia I Evolució*. 32:99–122.
- Company J, Pereda SX, Ruiz-Omenaca JJ. 2009. Nuevos restos fósiles del dinosaurio *Lirainosaurus* (Sauropoda, Titanosauria) en el Cretácico Superior (Campaniano-Maastrichtiano) de la Península Ibérica. *Ameghiniana*. 46:391–405.
- Cooper MR. 1981. The prosauropod dinosaur *Massospondylus Carinatus* Owen From Zimbabwe: its biology, mode of life and phylogenetic significance. *Occas Pap, Natl Mus*. 6:689–840.
- Cooper MR. 1984. A reassessment of *Vulcanodon karibaensis* raath (Dinosauria: saurischia) and the origin of the Sauropoda. *Palaeont Afr*. 25:203–231.
- Curry RK, Forster CA. 2004. The skull of *Rapetosaurus krausei* (Sauropoda: titanosauria) from the Late Cretaceous of Madagascar. *Journal of Vertebrate Paleontology*. 24:121–144.
- Deng SH, Lu YZ, Zhao Y, Fan R, Wang YD, Yang XJ, Li X, Sun BN. 2017. The Jurassic palaeoclimate regionalization and evolution of China. *Earth Sci Front*. 24:106–142.
- Dong ZM, Zhou SW, Zhang YZ. 1983. Dinosaurs from the jurassic of sichuan. In: Nanjing institute of geology and palaeontology and institute of vertebrate paleontology and paleoanthropology academia. editor. *Sinica Palaeontologica Sinica: Science Press*; p. 1–136.
- Fang XX, Long QQ, Lu LW, Zhang ZX, Pan SG, Wang WM, Li XK, Cheng ZW. 2000. Lower, middle, and upper jurassic subdivision in the lufeng region. Yunnan Province: 3th National Stratigraphical Congress of China; p. 208–214.
- Fang XX, Zhao XJ, Lu LW, Cheng ZW. 2004. Discovery of late jurassic *mamenchisaurus* in yunnan, Southwestern China. *Geol Bull China*. 23:1006–1009.
- Galton PM. 1984. Cranial anatomy of the prosauropod dinosaur *Plateosaurus* from the Knollenmergel (Middle Keuper, Upper Triassic) of Germany. *Geologica et Palaeontologica*. 18:139–171.
- Goloboff PA, Catalano SA. 2016. TNT version 1.5, including a full implementation of phylogenetic morphometrics. *Cladistics-The Int J Willi Hennig Soc*. 32:221–238.
- Harris JD. 2006a. Cranial osteology of *Suuwassea emilieae* (sauropoda: diplococoidea: flagellicaudata) from the upper jurassic morrison formation of montana, USA. *Journal of Vertebrate Paleontology*. 26:88–102.
- He XL, Li C, Cai KJ. 1988. The Middle Jurassic Dinosaur Fauna from Dashampu, Zigong, Sichuan: sauropod Dinosaurs, Volume 4, *Omeisaurus tianfuensis*. Chengdu: Sichuan Publishing House of Science and Technology; p. 143.
- He XL, Yang SH, Cai KJ, Li K, Liu ZW. 1996. A new species of sauropod, *Mamenchisaurus anyuesis* sp. nov. *Proc. 30th International Geological Congress*. 12: 83–86.
- Hou LH. 1976. New discovery of Sauropoda in Sichuan. *Vertebrata Palasiatica*. 14:160–165.
- Huang JD, You HL, Yang JT, Ren XX. 2014. A new sauropod dinosaur from the Middle Jurassic of Huangshan, Anhui Province. *Vertebrata Palasiatica*. 52:390–400.
- Huene FV. 1932. Die fossile Reptil-Ordnung Saurischia: ihre Entwicklung und Geschichte. Gebrüder Borntraeger. *Monographien Geologie und Palaontologie*. S1:1–361.
- Jain SL, Bandyopadhyad S. 1997. New titanosaurid (Dinosauria: sauropoda) from the Late Cretaceous of central India. *Journal of Vertebrate Paleontology*. 17:114–136.
- Jiang LL, Hu QZ, Zhu Q, Huang DZ, Wang DE. 2016. Late Mesozoic multi-stage structural deformations feature in the adjacent region among Anhui, Zhejiang, and Jiangxi Province. *Earth Sci Front*. 23:137–147.
- Jiang S, Li F, Peng GZ, Ye Y. 2011. A new species of *Omeisaurus* from the Middle Jurassic of Zigong, Sichuan. *Vertebrata Palasiatica*. 48:185–194.
- Knoll F, Ridgely RC, Ortega F, Sanz JL, Witmer LM. 2013. Neurocranial osteology and neuroanatomy of a Late Cretaceous titanosaurian sauropod from Spain (*Ampelosaurus* sp.). *PLoS One*. 8:e54991.
- Lehman TM, Coulson AB. 2002. A juvenile specimen of the Sauropod Dinosaur *Alamosaurus sanjuanensis* from the Upper Cretaceous of Big Bend National Park, Texas. *Paleontological Soc*. 76:156–172.
- Li K, Cai KJ. 1997. Classification and evolution of *Mamenchisaurus*. *J Chengdu Univ Technol*. 24:102–107.
- Li K, Yang CY, Liu J, Wang ZX. 2010. A new sauropod from Lower Jurassic of Huili, Sichuan, China. *Vertebrata Palasiatica*. 48:185–202.
- Lü JC, Li SX, Ji Q, Wang G, Zhang J, Dong Z. 2006. New eusauropod dinosaur from Yuanmou of Yunnan Province, China. *Acta Geologica Sinica*. 80:1–10.
- Lü JC, Li TG, Zhong SM, Ji Q, Li SX. 2008. A new mamenchisaurid dinosaur from the middle jurassic of yuanmou, yunnan province, china. *Acta Geologica Sinica*. 82:17–26.
- Mallison H. 2010a. The digital *Plateosaurus* I: body mass, mass distribution and posture assessed using CAD and CAE on a digitally mounted complete skeleton. *Palaeontologia Electronica*. 13:1–26.
- Mallison H. 2010b. The digital *Plateosaurus* II: an assessment of the range of motion of the limbs and vertebral column and of pervious reconstruction s using a digital skeletal mount. *Acta Palaeontol Pol*. 55:433–458.
- Mannion PD, Upchurch P, Barnes RN, Mateus O. 2013. Osteology of the Late Jurassic Portuguese sauropod dinosaur *Lusotitan atalaiensis* (Macronaria) and the evolutionary history of basal titanosauriforms. *Zool J Linn Soc*. 1–105.
- Mannion PD, Upchurch P, Carrano MT, Barrett PM. 2011. Testing the effect of the rock record on diversity: a multidisciplinary approach to elucidating the generic richness of sauropodomorph dinosaurs through time. *Biol Rev*. 86:157–181.
- Marsh OC. 1878. Principal characters of American Jurassic dinosaurs (Part 1). *Am J Sci*. 16:411–416.
- McIntosh JS. 1990. Sauropoda. In: Wieshampel DB, Dodson P, Omsolska H, editors. *The Dinosauria*. Berkeley: University of California Press; p. 345–407.
- McIntosh JS, Miller WE, Stadtman KL, Gillette DD. 1996. The Osteology of *Camarasaurus lewisi* (Jensen, 1988). *Byu Geol Stud*. 41:73–115.
- Monbaron M, Russell DA, Taquet P. 1999. *Atlasaurus imerakei* n. g., n. sp., a brachiosaurid-like sauropod from the Middle Jurassic of Morocco. *Comptes Rendus De l'Académie Des Sciences – Series IIA – Earth and Planetary Science*. 329:519–526.
- Ouyang H, Ye Y. 2002. The first mamenchisaurian skeleton with complete skull *Mamenchisaurus youngi*. Chengdu: Sichuan Publishing House of Science and Technology; p. 1–111.
- Powell JE. 1992. Osteologia de *Saltasaurus loricatus* (Sauropoda - Titanosauridae) del Cretácico Superior del Noroeste Argentino. In: Sanz JL, Buscalioni AD, editors. *Los Dinosaurios y Su Entorno Biotico: actas del Segundo Curso de Paleontología in Cuenca*. Instituto “Juan de Valdes”. Cuenca: Argentina; p. 165–230.
- Remes K. 2006. Revision of the Tendaguru sauropod dinosaur *Tornieria africana* (Fraas) and its relevance for sauropod paleobiogeography. *Journal of Vertebrate Paleontology*. 26:651–669.
- Remes K, Ortega F, Fierro I, Joger U, Kosma R, Marin Ferrer JM, Ide OA, Maga A. 2009. A new basal sauropod dinosaur from the middle jurassic of niger and the early evolution of sauropoda. *PLoS One*. 4:e6924.
- Ren XX, Huang JD, Hong TQ, Bi ZG. 2017. New research progress in Jurassic and Cretaceous stratigraphy of the Huangshan Area, Anhui Province. *J Stratigr*. 41:209–215.
- Ren XX, Huang JD, Liu J, Bi ZG. 2015. Late Jurassic volcanics discovered in the Tunxi Area of south Anhui. *Geol Anhui*. 25:241–244.
- Royo-Torres R, Cobos A, Alcalá L. 2006. A giant European dinosaur and a new sauropod clade. *Science*. 314:1925–1927.
- Royo-Torres R, Cobos A, Luque L, Aberasturi A, Espilez E, Fierro I, Gonzalez A, Mampel L, Alcalá L. 2009. High European sauropod dinosaur diversity during Jurassic-Cretaceous transition in Riodeva (Teruel, Spain). *Palaeontology*. 52:1009–1027.
- Royo-Torres R, Upchurch P. 2012. The cranial anatomy of the sauropod *Turiasaurus riodevensis* and implications for its phylogenetic relationships. *J Syst Palaeontology*. 10:553–583.
- Russell DA. 1993. The role of Central Asia in dinosaur biogeography. *Can J Earth Sci*. 30:2002–2012.
- Salgado L, Bonaparte JF. 1991. Un nuevo sauropodo Dicraeosauridae, *Amargasaurus cazaui* gen. et sp. nov., de la Formacion La Amarga,

- Neocomiano de la Provincia del Neuquén, Argentina. *Ameghiniana*. 28:333–346.
- Salgado L, Garrido A, Cocca SE, Cocca JR. 2004. Lower Cretaceous Rebbachisaurid Sauropods from Cerro Aguada Del Leon (Lohan Cura Formation), Neuquen Province, northwestern Patagonia, Argentina. *Journal of Vertebrate Paleontology*. 24:903–912.
- Sanz JL, Powell JE, Le Loeuff J, Martinez R, Pereda-Suberbiola X. 1999. Sauropod remains from the Upper Cretaceous of La ñ o (Northcentral Spain). titanosaur phylogenetic relationships. *Estudios del Museo de Ciencias Naturales de Alava*. 14:235–255.
- Seeley HG 1887. On the classification of the fossil animals commonly named Dinosauria. *Proceedings of the Royal Society of London*. 43: 165–171.
- Sereno PC, Wilson JA, Witmer LM, Whitlock J, Maga AA, Ide O, Rowe TA. 2007. Structural extremes in a Cretaceous dinosaur. *PLoS One*. 2:e1230.
- Suteethorn S, Le Loeuff J, Buffetaut E, Suteethorn V, Wongko K. 2012. First evidence of a mamenchisaurid dinosaur from the upper jurassic-lower cretaceous phu kratung formation of thailand. *Acta Palaeontologica*. 58:459–469.
- Tang F, Jing X, Kang X, Zhang G. 2001. *Omeisaurus maoianus*: A Complete Sauropod from Jingyuan, Sichuan. Beijing: China Ocean Press; p. 128.
- Upchurch P. 1995. The evolutionary history of sauropod dinosaurs. *Philosophical Trans Royal Soc London. Ser B: Biol Sci*. 349:365–390.
- Upchurch P. 1998. The phylogenetic relationships of sauropod dinosaurs. *Zool J Linn Soc*. 124:43–103.
- Upchurch P, Barrett PM, Dodson P. 2004. Sauropoda. In: Weishampel DB, Dodson P, Osmolska H, editors. *The Dinosauria*. Berkeley: University of California Press; p. 259–322.
- Wang LW. 2006. Hosting stratum and sedimentary facies of Mid-Jurassic dinosaur fossil relic in Jimu MT., Huangshan city. *Geol Anhui*. 16:161–164.
- Wang ZX, Li K, Liu J. 2003. *Mamenchisaurus* from Upper Jurassic of Jianyang, Sichuan. *J Chengdu Univ Technol*. 30:485–490.
- Whitlock JA, Harris JD. 2010. The dentary of *Suuwassea emilieae* (Sauropoda: diplodocoidea). *Journal of Vertebrate Paleontology*. 30:1637–1641.
- Wilson JA. 2002. Sauropod dinosaur phylogeny: critique and cladistics analysis. *Zool J Linn Soc*. 136:217–276.
- Wilson JA. 2005. Redescription of the Mongolian sauropod *Nemegtosaurus mongoliensis* Nowinski (Dinosauria: Saurischia) And Comments On Late Cretaceous sauropod diversity. *J Syst Palaeontology*. 3:283–318.
- Wilson JA, Sereno PC. 1998. Early evolution and higher-level phylogeny of sauropod dinosaurs. *Society Vertebrate Paleontology Memoir*. 5:1–68.
- Wilson JA, Upchurch P. 2003. A revision of *Titanosaurus lydekker* (Dinosauria-Sauropoda), the first dinosaur genus with a ‘Gondwanan’ distribution. *J Syst Palaeontology*. 1:125–160.
- Wilson JA, Upchurch P. 2009. Redescription and reassessment of the phylogenetic affinities of *Euhelopus zdanskyi* (Dinosauria: sauropoda) from the Early Cretaceous of China. *J Syst Palaeontology*. 7:199–239.
- Wu WH, Zhou CF, Oliver W, Sekiya T, Dong ZM. 2013. A new gigantic sauropod dinosaur from the Middle Jurassic of Shanshan, Xinjiang. *Global Geol*. 23:437–446.
- Xing LD, Miyashita T, Currie PJ, You HL, Zhang J, Dong ZM. 2015a. A new basal eusauropod from the Middle Jurassic of Yunnan, China, and faunal compositions and transitions of Asian sauropodomorph dinosaurs. *Acta Palaeontol Pol*. 60:145–154.
- Xing LD, Miyashita T, Zhang JP, Li DQ, Ye Y, Sekiya T, Wang FP, Currie PJ. 2015b. A new sauropod dinosaur from the Late Jurassic of China and the diversity, distribution, and relationships of mamenchisaurids. *Journal of Vertebrate Paleontology*. 35 (1):e889701.
- Young CC. 1939. On a new Sauropoda, with notes on other fragmentary reptiles from Szechuan. *Bull Geol Soc China*. 279–315.
- Young CC. 1954. On a new sauropod from Yiping, Szechuan, China. *Acta Palaeontologica Sinica*. 2:355–369.
- Young CC. 1958. New sauropods from China. *Vertebrata Palasiatica*. 2:1–29.
- Young CC, Chao XJ. 1972. *Mamenchisaurus hochuanensis*. In: Institute of Vertebrate Paleontology and Paleoanthropology Monograph Series I. editor. Beijing: Science Press; p. 1–30.
- Yu XQ, Wang DE. 2001. Jurassic-Cretaceous sequence stratigraphy of Huangshan Area in south Anhui. *Geoscience*. 15:27–34.
- Zhang YH, Li K, Zeng QH. 1998. A new species of sauropod dinosaur from the Upper Jurassic of Sichuan Basin, China. *J Chengdu Univ Technol*. 25:61–70.
- Zhao XJ. 1993. A new Middle Jurassic sauropod subfamily (Klamelisaurinae subfam. nov.) from Xinjiang Autonomous Region, China. *Vertebrata Palasiatica*. 31:132–138.